

Brown measures of sets of commuting operators in a type II_1 factor.

Hanne Schultz ^{*}[†]

Abstract

Using the spectral subspaces obtained in [HS], Brown's results (cf. [Bro]) on the Brown measure of an operator in a type II_1 factor $(\mathcal{M}, \tau)$ are generalized to finite sets of commuting operators in $\mathcal{M}$. It is shown that whenever $T_1, \dots, T_n \in \mathcal{M}$ are mutually commuting operators, there exists one and only one compactly supported Borel probability measure $\mu_{T_1, \dots, T_n}$ on $\mathbb{B}(\mathbb{C}^n)$ such that for all $\alpha_1, \dots, \alpha_n \in \mathbb{C}$,

$$\tau(\log |\alpha_1 T_1 + \dots + \alpha_n T_n - \mathbf{1}|) = \int_{\mathbb{C}^n} \log |\alpha_1 z_1 + \dots + \alpha_n z_n - 1| d\mu_{T_1, \dots, T_n}(z_1, \dots, z_n).$$

Moreover, for every polynomial q in n commuting variables, $\mu_{q(T_1, \dots, T_n)}$ is the push-forward measure of $\mu_{T_1, \dots, T_n}$ via the map $q : \mathbb{C}^n \rightarrow \mathbb{C}$.

In addition it is shown that, as in [HS], for every Borelset $B \subseteq \mathbb{C}^n$ there is a maximal closed $T_1, \dots, T_n$ -invariant subspace $\mathcal{K}$ affiliated with $\mathcal{M}$, such that $\mu_{T_1|_{\mathcal{K}}, \dots, T_n|_{\mathcal{K}}}$ is concentrated on B . Moreover, $\tau(P_{\mathcal{K}}) = \mu_{T_1, \dots, T_n}(B)$. This generalizes the main result from [HS] to n -tuples of commuting operators in $\mathcal{M}$.

1 Introduction.

In [Bro] Brown showed that for every operator T in a type II_1 factor $(\mathcal{M}, \tau)$ there is one and only one compactly supported Borel probability measure μ_T on $\mathbb{C}$, the *Brown measure* of T , such that for all $\lambda \in \mathbb{C}$,

$$\tau(\log |T - \lambda \mathbf{1}|) = \int_{\mathbb{C}} \log |z - \lambda| d\mu_T(z).$$

In [HS] it was shown that given $T \in \mathcal{M}$ and $B \in \mathbb{B}(\mathbb{C})$, there is a maximal closed T -invariant projection $P = P_T(B) \in \mathcal{M}$, such that the Brown measure of PTP (considered as an element of PMP) is concentrated on B . Moreover, $P_T(B)$ is hyperinvariant for T ,

$$\tau(P_T(B)) = \mu_T(B), \tag{1.1}$$

and the Brown measure of $P^{\perp}TP^{\perp}$ (considered as an element of $P^{\perp}\mathcal{M}P^{\perp}$) is concentrated on B^c .

In particular, if $S, T \in \mathcal{M}$ are commuting operators and $A, B \in \mathbb{B}(\mathbb{C})$, then $P_S(A) \wedge P_T(B)$ is S - and T -invariant. Thus, it is tempting to define a Brown measure for the pair (S, T) , $\mu_{S, T}$, by

$$\mu_{S, T}(A \times B) = \tau(P_S(A) \wedge P_T(B)), \quad A, B \in \mathbb{B}(\mathbb{C}). \tag{1.2}$$

^{*}As a student of the PhD-school OP-ALG-TOP-GEO the author is partially supported by the Danish Research Training Council.

[†]Partially supported by The Danish National Research Foundation.

In order to show that $\mu_{S,T}$ extends (uniquely) to a Borel probability measure on $\mathbb{C}^2$, we introduce the notion of an *idempotent valued measure* (cf. section 3), and for $T \in \mathcal{M}$ and $B \in \mathbb{B}(\mathbb{C})$ we define an unbounded idempotent affiliated with $\mathcal{M}$, $e_T(B)$, by $\mathcal{D}(e_T(B)) = \mathcal{K}_T(B) + \mathcal{K}_T(B^c)$ and

$$e_T(B)\xi = \begin{cases} \xi, & \xi \in \mathcal{K}_T(B) \\ 0, & \xi \in \mathcal{K}_T(B^c) \end{cases} \quad (1.3)$$

(cf. section 3). We prove that the map $B \mapsto e_T(B)$ is an idempotent valued measure and this enables us to show that $\mu_{S,T}$ given by (1.2) has an extension to a measure on $\mathbb{C}^2$ (cf. section 4). As in [HS] we also prove the existence of *spectral subspaces* for a set of commuting operators in $\mathcal{M}$. In the case of two commuting operators $S, T \in \mathcal{M}$, we show that for $B \in \mathbb{B}(\mathbb{C}^2)$ there is a maximal S - and T -invariant projection $P \in \mathcal{P}(\mathcal{M})$, such that $\mu_{S|P(\mathcal{H}), T|P(\mathcal{H})}$ (computed relative to PMP) is concentrated on B . In the case where $B = B_1 \times B_2$, $B_1, B_2 \in \mathbb{B}(\mathbb{C})$, P is simply the intersection $P_S(B_1) \wedge P_T(B_2)$ (cf. section 5).

Finally, in section 6 we show that $\mu_{S,T}$ has a characterization similar to the one given by Brown in [Bro]. That is, we show that $\mu_{S,T}$ is the unique compactly supported Borel probability measure on $\mathbb{B}(\mathbb{C}^2)$ such that for all $\alpha, \beta \in \mathbb{C}$,

$$\tau(\log |\alpha S + \beta T - \mathbf{1}|) = \int_{\mathbb{C}^2} \log |\alpha z + \beta w - 1| d\mu_{S,T}(z, w),$$

and there is a similar characterization of the Brown measure of an arbitrary finite set of commuting operators. In particular, $\mu_{\alpha S + \beta T}$ is the push-forward measure of $\mu_{S,T}$ via the map $(z, w) \mapsto \alpha z + \beta w$. In fact we can show that for every polynomial q in two commuting variables, $\mu_{q(S,T)}$ is the push-forward measure of $\mu_{S,T}$ via the map $q : \mathbb{C}^2 \rightarrow \mathbb{C}$. All this can (and will) be generalized to arbitrary finite sets of mutually commuting operators.

Section 2 is devoted to a summary of [Ne] in which we recall the definition of the measure topology on $\mathcal{M}$ and how the closure of $\mathcal{M}$ w.r.t. the measure topology may be identified with the set of closed, densely defined operators affiliated with $\mathcal{M}$. Afterwards we focus our attention on the set of unbounded idempotents affiliated with $\mathcal{M}$, $\mathcal{I}(\tilde{\mathcal{M}})$. We prove that these are in one-to-one correspondence with pairs of projections $P, Q \in \mathcal{M}$ with $P \wedge Q = 0$ and $P \vee Q = \mathbf{1}$. More precisely, when E is a closed, densely defined, unbounded operator affiliated with $\mathcal{M}$ with $\overline{E \cdot E} = E$, then P , the range projection of E , and Q , the projection onto the kernel of E , satisfy that $P \wedge Q = 0$ and $P \vee Q = \mathbf{1}$. Moreover, $\mathcal{D}(E) = P(\mathcal{H}) + Q(\mathcal{H})$, and E is given by

$$E\xi = \begin{cases} \xi, & \xi \in P(\mathcal{H}) \\ 0, & \xi \in Q(\mathcal{H}) \end{cases} \quad (1.4)$$

Vice versa, when $P, Q \in \mathcal{M}$ with $P \wedge Q = 0$ and $P \vee Q = \mathbf{1}$, then (1.4) defines a closed, densely defined, unbounded operator affiliated with $\mathcal{M}$ with $\overline{E \cdot E} = E$. In particular, (1.3) defines a closed, densely defined idempotent affiliated with $\mathcal{M}$. In section 2 it is also shown that $\mathcal{I}(\tilde{\mathcal{M}})$ is stable w.r.t. addition of countably many idempotents $(E_n)_{n=1}^\infty$ with $E_n E_m = E_m E_n = 0$, $n \neq m$. These are results which are needed in the succeeding sections.

Acknowledgements. Part of this work was carried out while I visited the UCLA Department of Mathematics. I want to thank the department, and especially the Operator Algebra Group, for their hospitality. I also thank my advisor, Uffe Haagerup, with whom I had enlightening discussions about this work.

2 Idempotents in $\tilde{\mathcal{M}}$.

We begin this section with a summary of E. Nelson's "Notes on non-commutative integration" [Ne]. We consider a finite von Neumann algebra $\mathcal{M}$ represented on a Hilbert space $\mathcal{H}$ and we fix a faithful, normal, tracial state τ on $\mathcal{M}$. We let $\mathcal{P}(\mathcal{M})$ denote the set of projections in $\mathcal{M}$.

Nelson defines the *measure topology* on $\mathcal{M}$ as follows: For $\varepsilon, \delta > 0$, let

$$N(\varepsilon, \delta) = \{T \in \mathcal{M} \mid \exists P \in \mathcal{P}(\mathcal{M}) : \|TP\| \leq \varepsilon, \tau(P^\perp) \leq \delta\}.$$

The measure topology on $\mathcal{M}$ is then the translation invariant topology on $\mathcal{M}$ for which the $N(\varepsilon, \delta)$'s form a fundamental system of neighborhoods of 0. $\tilde{\mathcal{M}}$ is the completion of $\mathcal{M}$ w.r.t. the measure topology.

Similarly, Nelson defines $\tilde{\mathcal{H}}$ to be the completion of $\mathcal{H}$ w.r.t. the translation invariant topology on $\mathcal{H}$ for which the sets

$$O(\varepsilon, \delta) = \{\xi \in \mathcal{H} \mid \exists P \in \mathcal{P}(\mathcal{M}) : \|P\xi\| \leq \varepsilon, \tau(P^\perp) \leq \delta\}$$

form a fundamental system of neighborhoods of 0. According to [Ne, Theorem 2], the natural mappings $\mathcal{M} \rightarrow \tilde{\mathcal{M}}$ and $\mathcal{H} \rightarrow \tilde{\mathcal{H}}$ are both injections.

2.1 Theorem. [Ne, Theorem 1] *The mappings*

$$\begin{aligned} \mathcal{M} \rightarrow \mathcal{M} & : T \mapsto T^*, \\ \mathcal{M} \times \mathcal{M} \rightarrow \mathcal{M} & : (S, T) \mapsto S + T, \\ \mathcal{M} \times \mathcal{M} \rightarrow \mathcal{M} & : (S, T) \mapsto S \cdot T, \\ \mathcal{H} \times \mathcal{H} \rightarrow \mathcal{H} & : (\xi, \eta) \mapsto \xi + \eta, \\ \mathcal{M} \times \mathcal{H} \rightarrow \mathcal{H} & : (T, \xi) \mapsto T\xi \end{aligned}$$

all have unique continuous extensions as mappings $\tilde{\mathcal{M}} \rightarrow \tilde{\mathcal{M}}$, $\tilde{\mathcal{M}} \times \tilde{\mathcal{M}} \rightarrow \tilde{\mathcal{M}}$, $\tilde{\mathcal{M}} \times \tilde{\mathcal{M}} \rightarrow \tilde{\mathcal{M}}$, $\tilde{\mathcal{H}} \times \tilde{\mathcal{H}} \rightarrow \tilde{\mathcal{H}}$ and $\tilde{\mathcal{M}} \times \tilde{\mathcal{H}} \rightarrow \tilde{\mathcal{H}}$, respectively. In particular, $\tilde{\mathcal{H}}$ is a complex vector space and $\tilde{\mathcal{M}}$ is a complex $$ -algebra with a continuous representation on $\tilde{\mathcal{H}}$.*

For $x \in \tilde{\mathcal{M}}$ define

$$\mathcal{D}(M_x) = \{\xi \in \mathcal{H} \mid x\xi \in \mathcal{H}\} \tag{2.1}$$

and define $M_x : \mathcal{D}(M_x) \rightarrow \mathcal{H}$ by

$$M_x \xi = x\xi, \quad \xi \in \mathcal{D}(M_x). \tag{2.2}$$

Recall that a (not necessarily bounded or everywhere defined) operator A on $\mathcal{H}$ is said to be *affiliated* with $\mathcal{M}$ if $AU = UA$ for every unitary $U \in \mathcal{M}'$.

2.2 Theorem. [Ne, Theorem 4] *For every $x \in \tilde{\mathcal{M}}$, M_x is a closed, densely defined operator affiliated with $\mathcal{M}$, and*

$$M_x^* = M_{x^*}. \tag{2.3}$$

Moreover, for $x, y \in \tilde{\mathcal{M}}$,

$$M_{x+y} = \overline{M_x + M_y}, \tag{2.4}$$

$$M_{x \cdot y} = \overline{M_x \cdot M_y}, \tag{2.5}$$

where $\overline{A}$ denotes the closure of a closable operator A .

A closed, densely defined operator A on $\mathcal{H}$ has a polar decomposition $A = V|A|$, and if A is affiliated with $\mathcal{M}$, then $V \in \mathcal{M}$ and all the spectral projections $(E_{|A|}([0, t]))_{t>0}$ belong to $\mathcal{M}$. Put

$$A_n = V \int_0^n t \, dE_{|A|}(t).$$

Assuming that A is affiliated with $\mathcal{M}$, we get that $(A_n)_{n=1}^\infty$ is a Cauchy sequence w.r.t. the measure topology. Indeed,

$$(A_{n+k} - A_n) \cdot E_{|A|}([0, n]) = 0,$$

and $\tau(E_{|A|}([n, \infty))) \rightarrow 0$ as $n \rightarrow \infty$. Hence, there exists $a \in \tilde{\mathcal{M}}$ s.t. $A_n \rightarrow a$ in measure, and according to [Ne, Theorem 3], $A = M_a$. It follows that every closed, densely defined operator A affiliated with $\mathcal{M}$ is of the form $A = M_a$ for some $a \in \tilde{\mathcal{M}}$, and this a is uniquely determined. Indeed, if $M_a = M_b$, then a and b agree on $\mathcal{D}(M_a)$ which is dense in $\mathcal{H}$ w.r.t. the norm topology and hence dense in $\tilde{\mathcal{H}}$ w.r.t. the measure topology. Since the representation of $\tilde{\mathcal{M}}$ on $\mathcal{H}$ is continuous, it follows that a and b agree on all of $\tilde{\mathcal{H}}$. By the same argument, if S and T are closed, densely defined operators affiliated with $\mathcal{M}$ which agree on a dense subset of $\mathcal{H}$, then $S = T$.

In summary, $\tilde{\mathcal{M}}$ is the completion of $\mathcal{M}$ w.r.t. the measure topology but it may also be viewed as the set of closed, densely defined operators affiliated with $\mathcal{M}$. In particular, if S and T belong to the latter, then Theorem 2.2 tells us that S^* , $\overline{S + T}$ and $\overline{S \cdot T}$ are also closed, densely defined and affiliated with $\mathcal{M}$.

2.3 Notation. In what follows we shall identify $\tilde{\mathcal{M}}$ with the set of closed, densely defined operators affiliated with $\mathcal{M}$, but whenever necessary, we will specify which one of the two pictures mentioned above we are using. In general, lower case letters will represent elements of the completion of $\mathcal{M}$ w.r.t. the measure topology, whereas upper case letters will represent closed, densely defined operators affiliated with $\mathcal{M}$. For $x \in \tilde{\mathcal{M}}$ we will denote by $\ker(x)$, $\text{range}(x)$ and $\text{supp}(x)$ the kernel of M_x , the range of M_x and the support projection of M_x , respectively.

In the rest of this section we will study the set of idempotent elements in $\tilde{\mathcal{M}}$,

$$\mathcal{I}(\tilde{\mathcal{M}}) = \{e \in \tilde{\mathcal{M}} \mid e \cdot e = e\}. \quad (2.6)$$

Alternatively, if $\tilde{\mathcal{M}}$ is viewed as the set of closed, densely defined operators affiliated with $\mathcal{M}$, then $\mathcal{I}(\tilde{\mathcal{M}})$ is the subset of operators E fulfilling that $\overline{E \cdot E} = E$.

The following proposition shows that $\mathcal{I}(\tilde{\mathcal{M}})$ is in one-to-one correspondence with the set of pairs of projections $P, Q \in \mathcal{M}$ such that $P \wedge Q = 0$ and $P \vee Q = \mathbf{1}$.

2.4 Proposition. *Let $E \in \mathcal{I}(\tilde{\mathcal{M}})$. Then the range of E , $\text{range}(E)$, and the kernel of E , $\ker(E)$ (which is the range of $\mathbf{1} - E$), are closed subspaces of $\mathcal{H}$, with*

$$\text{range}(E) \cap \ker(E) = \{0\} \quad (2.7)$$

and

$$\overline{\text{range}(E) + \ker(E)} = \mathcal{H}. \quad (2.8)$$

Moreover, $\mathcal{D}(E) = \text{range}(E) + \ker(E)$. Conversely, if $P, Q \in \mathcal{M}$ are projections with $P \wedge Q = 0$ and $P \vee Q = \mathbf{1}$, then there is a unique idempotent $E \in \tilde{\mathcal{M}}$ with $\mathcal{D}(E) = P(\mathcal{H}) + Q(\mathcal{H})$, $\text{range}(E) = P(\mathcal{H})$ and $\ker(E) = Q(\mathcal{H})$, and E is determined by

$$E\xi = \begin{cases} \xi, & \xi \in P(\mathcal{H}), \\ 0, & \xi \in Q(\mathcal{H}). \end{cases} \quad (2.9)$$

Proof. Write $E = M_e$ for a (uniquely determined) element $e \in \tilde{\mathcal{M}}$ with $e \cdot e = e$ and recall from (2.1) that

$$\mathcal{D}(E) = \{\xi \in \mathcal{H} \mid e\xi \in \mathcal{H}\}.$$

Let $E \cdot E$ denote the densely defined operator on $\mathcal{H}$ obtained by composing E with itself. Then

$$\begin{aligned} \mathcal{D}(E \cdot E) &= \{\xi \in \mathcal{D}(E) \mid E\xi \in \mathcal{D}(E)\} \\ &= \{\xi \in \mathcal{D}(E) \mid eE\xi \in \mathcal{H}\} \\ &= \{\xi \in \mathcal{D}(E) \mid e^2\xi \in \mathcal{H}\}. \end{aligned}$$

But $e^2 = e$ as everywhere defined operators on $\tilde{\mathcal{H}}$. Hence, for $\xi \in \mathcal{D}(E)$,

$$e^2\xi = e\xi = E\xi,$$

and it follows that $\mathcal{D}(E \cdot E) = \mathcal{D}(E)$ and that $E \cdot E = E$ (without taking closure). In particular, $\text{range}(E) \subseteq \mathcal{D}(E)$. Moreover, since $E \cdot E = E$,

$$\text{range}(E) = \{\xi \in \mathcal{D}(E) \mid E\xi = \xi\} = \ker(\mathbf{1} - E).$$

According to [KR, Exercise 2.8.45], the kernel of a closed operator is closed. Hence, $\ker(E)$ and $\text{range}(E) = \ker(\mathbf{1} - E)$ are closed. Moreover,

$$\text{range}(E) \cap \ker(\mathbf{1} - E) = \ker(\mathbf{1} - E) \cap \ker(E) = \{0\}.$$

Clearly, $\text{range}(E) + \ker(E) \subseteq \mathcal{D}(E)$, and since

$$\xi = E\xi + (\mathbf{1} - E)\xi, \quad \xi \in \mathcal{D}(E),$$

the converse inclusion also holds. That is,

$$\mathcal{D}(E) = \text{range}(E) + \ker(E)$$

Let P and Q denote the projections onto $\text{range}(E)$ and $\ker(E)$, respectively. Then by the above, $P \wedge Q = 0$ and $P \vee Q = \mathbf{1}$, and E is the operator on $\mathcal{D}(E) = P(\mathcal{H}) + Q(\mathcal{H})$ determined by (2.9). In particular, E is uniquely determined by its range and its kernel.

Conversely, assume that P and Q are projections in $\mathcal{M}$, such that $P \wedge Q = 0$ and $P \vee Q = \mathbf{1}$, and let E be the operator on $\mathcal{D}(E) = P(\mathcal{H}) + Q(\mathcal{H})$ determined by (2.9). Then clearly, $E \cdot E = E$ (without taking closure), $\text{range}(E) = P(\mathcal{H})$, and $\ker(E) = Q(\mathcal{H})$. Moreover, the graph of E is given by

$$\begin{aligned} \mathcal{G}(E) &= \{(\xi + \eta, \xi) \mid \xi \in P(\mathcal{H}), \eta \in Q(\mathcal{H})\} \\ &= \{(u, v) \in \mathcal{H} \times \mathcal{H} \mid u - v \in Q(\mathcal{H}), v \in P(\mathcal{H})\}, \end{aligned}$$

which is clearly a closed subspace of $\mathcal{H} \times \mathcal{H}$. Hence, $E \in \mathcal{I}(\tilde{\mathcal{M}})$. ■

2.5 Definition. We define $\text{tr} : \mathcal{I}(\tilde{\mathcal{M}}) \rightarrow [0, 1]$ by

$$\text{tr}(e) = \tau(\text{supp}(e)), \quad e \in \mathcal{I}(\tilde{\mathcal{M}}), \tag{2.10}$$

where $\text{supp}(e) \in \mathcal{P}(\mathcal{M})$ denotes the support projection of M_e .

2.6 Remark. For every $x \in \tilde{\mathcal{M}}$, $\text{supp}(x) \sim P_{\text{range}(x)}$ so one also has that for $e \in \mathcal{J}(\tilde{\mathcal{M}})$,

$$\text{tr}(e) = \tau(P_{\text{range}(e)}). \quad (2.11)$$

Throughout the paper we will, without further mentioning, make use of this identity.

2.7 Proposition. Let $e_1, \dots, e_n \in \mathcal{J}(\tilde{\mathcal{M}})$ (seen as everywhere defined operators on $\tilde{\mathcal{H}}$) with $e_i e_j = 0$ when $i \neq j$. Then $e_1 + \dots + e_n \in \mathcal{J}(\tilde{\mathcal{M}})$, and

- (a) $\ker(e_1 + \dots + e_n) = \bigcap_{i=1}^n \ker(e_i)$,
- (b) $\text{supp}(e_1 + \dots + e_n) = \bigvee_{i=1}^n \text{supp}(e_i)$,
- (c) $\text{range}(e_1 + \dots + e_n) = \bigvee_{i=1}^n \text{range}(e_i)$,
- (d) $\text{tr}(e_1 + \dots + e_n) = \sum_{i=1}^n \text{tr}(e_i)$.

Proof. It suffices to consider the case $n = 2$. The general case follows by induction over $n \in \mathbb{N}$. If $e_1, e_2 \in \mathcal{J}(\tilde{\mathcal{M}})$ with $e_1 e_2 = e_2 e_1 = 0$, then obviously, $e_1 + e_2 \in \mathcal{J}(\tilde{\mathcal{M}})$.

(a) Clearly, $\ker(e_1) \cap \ker(e_2) \subseteq \ker(e_1 + e_2)$. On the other hand, if $\xi \in \ker(e_1 + e_2)$, then $e_i \xi = e_i(e_1 + e_2)\xi = 0$, ($i = 1, 2$), whence $\ker(e_1 + e_2) \subseteq \ker(e_1) \cap \ker(e_2)$.

(b) Since $\text{supp}(e_1 + e_2)$ is the projection onto $[\ker(e_1 + e_2)]^\perp$, (b) follows from (a).

(c) For a closed, densely defined operator S affiliated with $\mathcal{M}$, $\overline{\text{range}(S)} = \ker(S^*)^\perp$. Using that every idempotent has closed range (cf. Proposition 2.4) and applying (a) to $e_1^* + \dots + e_n^*$, we find that

$$\begin{aligned} \text{range}(e_1 + \dots + e_n) &= \overline{\text{range}(e_1 + \dots + e_n)} \\ &= [\ker(e_1^* + \dots + e_n^*)]^\perp \\ &= \bigvee_{i=1}^n \ker(e_i^*)^\perp \\ &= \bigvee_{i=1}^n \text{range}(e_i). \end{aligned}$$

(d) For $i = 1, 2$ put $P_i = P_{\text{range}(e_i)}$. Since $e_1 e_2 = e_2 e_1 = 0$, for $\xi \in P_1(\mathcal{H}) \cap P_2(\mathcal{H})$ we have that

$$\xi = e_1 \xi = e_2 e_1 \xi = 0.$$

Then by Kaplansky's formula (cf. [KR, Theorem 6.1.7]),

$$P_1 \vee P_2 - P_1 \sim P_2 - P_1 \wedge P_2 = P_2,$$

so that

$$\tau(P_1 \vee P_2) = \tau(P_1) + \tau(P_2).$$

According to (c), $P_1 \vee P_2$ is the projection onto $\text{range}(e_1 + e_2)$, and then by Remark 2.6,

$$\text{tr}(e_1 + e_2) = \text{tr}(e_1) + \text{tr}(e_2). \quad \blacksquare$$

2.8 Proposition. Let $(e_n)_{n=1}^\infty$ be a sequence of idempotents in $\tilde{\mathcal{M}}$ with $e_n e_m = e_m e_n = 0$ when $n \neq m$. Then there is an idempotent in $\tilde{\mathcal{M}}$, which we denote by $\sum_{n=1}^\infty e_n$, such that $\sum_{n=1}^N e_n \rightarrow \sum_{n=1}^\infty e_n$ in measure as $N \rightarrow \infty$. Moreover, $\text{supp}(\sum_{n=1}^N e_n) \nearrow \text{supp}(\sum_{n=1}^\infty e_n)$, whence

$$\text{supp}\left(\sum_{n=1}^\infty e_n\right) = \bigvee_{n=1}^\infty \text{supp}(e_n), \quad (2.12)$$

and

$$\text{tr}\left(\sum_{n=1}^\infty e_n\right) = \sum_{n=1}^\infty \text{tr}(e_n). \quad (2.13)$$

Also,

$$\text{range}\left(\sum_{n=1}^\infty e_n\right) = \bigvee_{n=1}^\infty \text{range}(e_n). \quad (2.14)$$

Proof. Let

$$f_n = \sum_{k=1}^n e_k, \quad n \in \mathbb{N}. \quad (2.15)$$

Then, according to Proposition 2.7, $\text{supp}(f_n) = \bigvee_{k=1}^n \text{supp}(e_k)$, and $\text{tr}(f_n) = \sum_{k=1}^n \text{tr}(e_k)$. We prove that $(f_n)_{n=1}^\infty$ is a Cauchy sequence in $\tilde{\mathcal{M}}$. For $n, k \in \mathbb{N}$, let

$$P_{n,k} = \text{supp}(f_{n+k} - f_n) = \bigvee_{l=1}^k \text{supp}(e_{n+l}). \quad (2.16)$$

Then

$$(f_{n+k} - f_n)P_{n,k}^\perp = 0, \quad (2.17)$$

so for every $\varepsilon > 0$,

$$f_{n+k} - f_n \in \mathcal{N}(\varepsilon, \tau(P_{n,k})) = \mathcal{N}\left(\varepsilon, \sum_{l=1}^k \text{tr}(e_{n+l})\right). \quad (2.18)$$

Now, $\sum_{k=1}^n \text{tr}(e_k) = \text{tr}(f_n) \leq 1$, so for arbitrary $\delta > 0$ there is an $n_0 \in \mathbb{N}$ such that

$$\sum_{k=n_0}^\infty \text{tr}(e_k) \leq \delta. \quad (2.19)$$

It follows from (2.18) and (2.19) that when $n \geq n_0$ and $k \geq 1$, then $f_{n+k} - f_n \in \mathcal{N}(\varepsilon, \delta)$. Thus, $(f_n)_{n=1}^\infty$ is a Cauchy sequence in $\tilde{\mathcal{M}}$. Put

$$e = \lim_{n \rightarrow \infty} f_n \in \tilde{\mathcal{M}}. \quad (2.20)$$

For all $k, n \in \mathbb{N}$, $f_{n+k}f_n = f_n f_{n+k} = f_n$, and hence

$$ef_n = f_n e = f_n, \quad (2.21)$$

so that $e \cdot e = e$.

Let $P_n = \text{supp}(f_n) = \bigvee_{k=1}^n \text{supp}(e_k)$. Then $P_n \nearrow P := \bigvee_{k=1}^{\infty} \text{supp}(e_k)$ as $n \rightarrow \infty$, and

$$\begin{aligned} \sum_{k=1}^{\infty} \text{tr}(e_k) &= \lim_{n \rightarrow \infty} \sum_{k=1}^n \text{tr}(e_k) \\ &= \lim_{n \rightarrow \infty} \text{tr}(f_n) \\ &= \lim_{n \rightarrow \infty} \tau(P_n) \\ &= \text{tr}(P). \end{aligned}$$

It follows from (2.21) that for every $n \in \mathbb{N}$, $P_n \leq \text{supp}(e)$. Hence $P \leq \text{supp}(e)$. On the other hand, for every $n \in \mathbb{N}$, $f_n(\mathbf{1} - P_n) = 0$, so

$$e(\mathbf{1} - P) = \lim_{n \rightarrow \infty} [f_n(\mathbf{1} - P_n)] = 0 \quad (2.22)$$

(the limit refers to the measure topology). Thus, $\text{supp}(e) \leq P$, and we have shown that $\text{supp}(e) = P = \bigvee_{k=1}^{\infty} \text{supp}(e_k)$ and that (2.13) holds.

In order to prove (2.14), let $\xi \in \text{range}(e_m)$. Then $e_m \xi = \xi$ and

$$\left(\sum_{n=1}^{\infty} e_n \right) \xi = e_m \xi = \xi.$$

Thus,

$$\text{range} \left(\sum_{n=1}^{\infty} e_n \right) \supseteq \bigvee_{n=1}^{\infty} \text{range}(e_n). \quad (2.23)$$

On the other hand, we know that $\text{range} \left(\sum_{n=1}^{\infty} e_n \right) \subseteq \mathcal{H}$, and since $\text{range} \left(\sum_{n=1}^N e_n \right) \subseteq \bigvee_{n=1}^{\infty} \text{range}(e_n)$ for all $N \in \mathbb{N}$, we also have that

$$\text{range} \left(\sum_{n=1}^{\infty} e_n \right) \subseteq \left(\bigvee_{n=1}^{\infty} \text{range}(e_n) \right)^{\sim},$$

where $\sim$ denotes closure w.r.t. the measure topology. Intersecting by $\mathcal{H}$ on both sides of the inclusion, we get that

$$\text{range} \left(\sum_{n=1}^{\infty} e_n \right) \subseteq \bigvee_{n=1}^{\infty} \text{range}(e_n). \quad (2.24)$$

This proves (2.14). $\blacksquare$

We shall make use of the following theorem from [A]. For a published proof of it, we refer the reader to [Aa].

2.9 Theorem. [A] Let E and F be (not necessarily closed) subspaces of $\mathcal{H}$ which are affiliated with $\mathcal{M}$.¹ Then $E \cap F$ is affiliated with $\mathcal{A}$, and

$$\overline{E \cap F} = \overline{E} \cap \overline{F}. \quad (2.25)$$

¹A subspace E of $\mathcal{H}$ is said to be *affiliated with* $\mathcal{M}$ if for all $T \in \mathcal{M}'$, $T(E) \subseteq E$. Note that if E is affiliated with $\mathcal{M}$, then the projection onto $\overline{E}$ belongs to $\mathcal{M}$, and if E is *closed*, then this is a necessary and sufficient condition for E to be affiliated with $\mathcal{M}$.

2.10 Lemma. Consider idempotents $e, f \in \tilde{\mathcal{M}}$. Let $P = P_{\text{range}(e)}$, $Q = P_{\text{range}(\mathbf{1}-e)}$, $R = P_{\text{range}(f)}$ and $S = P_{\text{range}(\mathbf{1}-f)}$. Then $ef = fe$ if and only if

$$(P \wedge R) \vee (P \wedge S) \vee (Q \wedge R) \vee (Q \wedge S) = \mathbf{1}. \quad (2.26)$$

Proof. Clearly,

$$\mathbf{1} = ef + e(\mathbf{1} - f) + (\mathbf{1} - e)f + (\mathbf{1} - e)(\mathbf{1} - f). \quad (2.27)$$

Suppose that $ef = fe$, and let $g_1 = ef$, $g_2 = e(\mathbf{1} - f)$, $g_3 = (\mathbf{1} - e)f$ and $g_4 = (\mathbf{1} - e)(\mathbf{1} - f)$. Then $g_1, \dots, g_4$ are idempotents with support projections P_1, P_2, P_3 and P_4 , respectively, such that

$$\left(\bigvee_{i=1}^4 P_i \right) (\mathcal{H}) = \mathcal{H}. \quad (2.28)$$

Moreover, $P_1 \leq P \wedge R$, $P_2 \leq P \wedge S$, $P_3 \leq Q \wedge R$ and $P_4 \leq Q \wedge S$. This shows that (2.26) holds.

On the other hand, assume that (2.26) holds. According to (2.26) and Theorem 2.9,

$$\mathcal{H}_0 := \mathcal{D}(ef) \cap \mathcal{D}(fe) \cap [(P \wedge R)(\mathcal{H}) + (P \wedge S)(\mathcal{H}) + (Q \wedge R)(\mathcal{H}) + (Q \wedge S)(\mathcal{H})]$$

is dense in $\mathcal{H}$ so it suffices to prove that ef and fe agree on $\mathcal{H}_0$. To see this, let $\xi \in \mathcal{D}(ef) \cap \mathcal{D}(fe) \cap (P \wedge R)(\mathcal{H})$. Then

$$ef\xi = e\xi = \xi = f\xi = fe\xi, \quad (2.29)$$

and similarly, when $\xi \in \mathcal{D}(ef) \cap \mathcal{D}(fe) \cap (P \wedge S)(\mathcal{H})$, $\xi \in \mathcal{D}(ef) \cap \mathcal{D}(fe) \cap (Q \wedge R)(\mathcal{H})$ or $\xi \in \mathcal{D}(ef) \cap \mathcal{D}(fe) \cap (Q \wedge S)(\mathcal{H})$. Thus, ef agrees with fe on $\mathcal{H}_0$. ■

3 An idempotent valued measure associated with $T \in \mathcal{M}$.

As in the previous section, consider a finite von Neumann algebra $\mathcal{M}$ with a faithful, normal, tracial state τ . Inspired by the notion of a spectral measure we make the following definition:

3.1 Definition. Let $(X, \mathcal{F})$ denote a measurable space. An *idempotent valued measure* on $(X, \mathcal{F})$ (with values in $\tilde{\mathcal{M}}$) is a map e from $\mathcal{F}$ into $\mathcal{J}(\tilde{\mathcal{M}})$ such that

- (i) $e(X) = \mathbf{1}$,
- (ii) $e(F_1)e(F_2) = e(F_2)e(F_1) = 0$ when $F_1, F_2 \in \mathcal{F}$ with $F_1 \cap F_2 = \emptyset$,
- (iii) when $(F_n)_{n=1}^\infty$ is a sequence of mutually disjoint sets from $\mathcal{F}$, then $\sum_{n=1}^N e(F_n)$ converges in measure as $N \rightarrow \infty$ to $e\left(\bigcup_{n=1}^\infty F_n\right)$, i.e.

$$e\left(\bigcup_{n=1}^\infty F_n\right) = \sum_{n=1}^\infty e(F_n).$$

Note that because of (ii) and Proposition 2.8, the limit in (iii) actually exists.

From now on we will assume that $\mathcal{M}$ is in fact a type II_1 factor. Recall from [HS] that for $T \in \mathcal{M}$ and $B \subseteq \mathbb{C}$ a Borel set there is a maximal T -invariant projection $P = P_T(B) \in \mathcal{M}$, such that the

Brown measure of PTP (considered as an element of PMP) is concentrated on B . Moreover, $P_T(B)$ is hyperinvariant for T ,

$$\tau(P_T(B)) = \mu_T(B), \quad (3.1)$$

and the Brown measure of $P^\perp TP^\perp$ (considered as an element of $P^\perp \mathcal{M} P^\perp$) is concentrated on B^c . We let $\mathcal{K}_T(B)$ denote the range of $P_T(B)$. Then the aim of this section is to prove:

3.2 Theorem. *Let $T \in \mathcal{M}$, and for $B \in \mathbb{B}(\mathbb{C})$, let $e_T(B)$ with $\mathcal{D}(e_T(B)) = \mathcal{K}_T(B) + \mathcal{K}_T(B^c)$ be given by*

$$e_T(B)\xi = \begin{cases} \xi, & \xi \in \mathcal{K}_T(B) \\ 0, & \xi \in \mathcal{K}_T(B^c) \end{cases} \quad (3.2)$$

Then $e_T(B) \in \mathcal{J}(\tilde{\mathcal{M}})$, and $B \mapsto e_T(B)$ is an idempotent valued measure.

The proof of this theorem uses various results which we state and prove below. The first one of these is a lemma which we proved in [HS], but for the sake of completeness we give the proof here as well.

3.3 Lemma. *Let $T \in \mathcal{M}$, and let $P \in \mathcal{M}$ be a non-zero, T -invariant projection. Then for every $B \in \mathbb{B}(\mathbb{C})$,*

$$\mathcal{K}_{T|_{P(\mathcal{H})}}(B) = \mathcal{K}_T(B) \cap P(\mathcal{H}), \quad (3.3)$$

where $T|_{P(\mathcal{H})}$ is considered as an element of the type II_1 factor PMP .

Proof. Let $Q \in PMP$ denote the projection onto $\mathcal{K}_{T|_{P(\mathcal{H})}}(B)$, and let $R = P_T(B) \wedge P$. We will prove that $Q \leq R$ and $R \leq Q$.

Clearly, $Q \leq P$. In order to see that $Q \leq P_T(B)$, recall that $P_T(B)$ is maximal w.r.t. the properties

- (i) $P_T(B)TP_T(B) = TP_T(B)$,
- (ii) $\mu_{P_T(B)TP_T(B)}$ (computed relative to $P_T(B)\mathcal{M}P_T(B)$) is concentrated on B .

Since

$$QTQ = QTPQ = TPQ = TQ, \quad (3.4)$$

and $\mu_{QTQ} = \mu_{QTPQ}$ (computed relative to QM) is concentrated on B , we get that $Q \leq P_T(B)$, and hence $Q \leq R$.

Similarly, to prove that $R \leq Q$, we must show that

- (i') $RTPR = TPR$, i.e. $RTR = TR$,
- (ii') $\mu_{RTPR} = \mu_{RTR}$ (computed relative to RM) is concentrated on B .

Note that if $P_T(B) = 0$, then $R \leq Q$, so we may assume that $P_T(B) \neq 0$. (i') holds, because $R(\mathcal{H}) = P(\mathcal{H}) \cap P_T(B)(\mathcal{H})$ is T -invariant when $P(\mathcal{H})$ and $P_T(B)(\mathcal{H})$ are T -invariant. In order to prove (ii'), at first note that $R(\mathcal{H})$ is $TP_T(B)$ -invariant. Hence

$$\mu_{TP_T(B)} = \tau_1(R) \cdot \mu_{RTR} + \tau_1(R^\perp) \cdot \mu_{R^\perp TR^\perp}, \quad (3.5)$$

where $\tau_1 = \frac{1}{\tau(P_T(B))} \cdot \tau|_{P_T(B)\mathcal{M}P_T(B)}$. It follows that

$$\tau_1(R) \cdot \mu_{RTR}(B^c) \leq \mu_{TP_T(B)}(B^c) = 0, \quad (3.6)$$

and thus, if $R \neq 0$, then $\mu_{RTR}(B^c) = 0$, and (ii') holds. If $R = 0$, then $R \leq Q$ is trivially fulfilled. ■

3.4 Proposition. For every Borel set $B \subseteq \mathbb{C}$,

$$\mathcal{K}_T(B) = \mathcal{K}_{T^*}((B^c)^*)^\perp, \quad (3.7)$$

where $A^* := \{\bar{z} \mid z \in A\}$ for $A \subseteq \mathbb{C}$. Moreover, for all Borel sets $A, B \subseteq \mathbb{C}$,

$$\mathcal{K}_T(A) \cap \mathcal{K}_T(B) = \mathcal{K}_T(A \cap B), \quad (3.8)$$

and

$$\mathcal{K}_T(A \cup B) = \overline{\mathcal{K}_T(A) + \mathcal{K}_T(B)}. \quad (3.9)$$

Proof. Let $B \in \mathbb{B}(\mathbb{C})$ and let $P = P_T(B)$. Then $P^\perp$ is T^* -invariant, and

$$\mu_{P^\perp T^* P^\perp}(B^*) = \mu_{(P^\perp T^* P^\perp)^*}(B) = \mu_{P^\perp T P^\perp}(B) = 0 \quad (3.10)$$

(recall that $\mu_{P^\perp T P^\perp}$ is concentrated on B^c). Thus, $\mu_{P^\perp T^* P^\perp}$ is concentrated on $\mathbb{C} \setminus B^*$, and maximality of $P_{T^*}(\mathbb{C} \setminus B^*)$ implies that

$$P_T(B)^\perp = P^\perp \leq P_{T^*}(\mathbb{C} \setminus B^*). \quad (3.11)$$

Since

$$\begin{aligned} \tau(P_{T^*}(\mathbb{C} \setminus B^*)) &= \mu_{T^*}(\mathbb{C} \setminus B^*) \\ &= 1 - \mu_{T^*}(B^*) \\ &= 1 - \mu_T(B) \\ &= \tau(P_T(B)^\perp), \end{aligned}$$

we get from (3.11) that $P_T(B)^\perp = P_{T^*}(\mathbb{C} \setminus B^*)$.

Next, let $A, B \in \mathbb{B}(\mathbb{C})$. By maximality of $P_T(A)$ and $P_T(B)$, $P_T(A \cap B) \leq P_T(A) \wedge P_T(B)$, so $\supseteq$ holds in (3.8). We let $\mathcal{K} := \mathcal{K}_T(A) \cap \mathcal{K}_T(B)$, and we let Q denote the projection onto $\mathcal{K}$. Then, according to Lemma 3.3,

$$\begin{aligned} \mathcal{K} &= \mathcal{K}_{T|_{\mathcal{K}_T(A)}}(B) \\ &= \mathcal{K}_{T|_{\mathcal{K}_T(B)}}(A), \end{aligned}$$

proving that μ_{QTQ} is concentrated on A and on B and therefore on $A \cap B$. Consequently, $Q \leq P_T(A \cap B)$, so $\subseteq$ also holds in (3.8).

Finally, we infer from (3.7) and (3.8) that

$$\begin{aligned} \mathcal{K}_T(A \cup B) &= \mathcal{K}_T((A^c \cap B^c)^c) \\ &= \mathcal{K}_{T^*}((A^c \cap B^c)^*)^\perp \\ &= \mathcal{K}_{T^*}((A^c)^* \cap (B^c)^*)^\perp \\ &= [\mathcal{K}_{T^*}((A^c)^*) \cap \mathcal{K}_{T^*}((B^c)^*)]^\perp \\ &= \overline{\mathcal{K}_{T^*}((A^c)^*)^\perp + \mathcal{K}_{T^*}((B^c)^*)^\perp} \\ &= \overline{\mathcal{K}_T(A) + \mathcal{K}_T(B)}. \quad \blacksquare \end{aligned}$$

It follows from Proposition 3.4 and Proposition 2.4 that for $B \in \mathbb{B}(\mathbb{C})$, $e_T(B)$ given by (3.2) belongs to $\mathcal{J}(\tilde{\mathcal{M}})$, as stated in Theorem 3.2.

3.5 Lemma. *Let $(x_n)_{n=1}^\infty$ be a sequence in $\tilde{\mathcal{M}}$, and suppose $\tau(\text{supp}(x_n)) \rightarrow 0$ as $n \rightarrow \infty$. Then $x_n \rightarrow 0$ in the measure topology.*

Proof. This is standard. $\blacksquare$

If $S \in \mathcal{M}$ commutes with $T \in \mathcal{M}$, then for every $B \in \mathbb{B}(\mathbb{C})$, $\mathcal{K}_T(B)$ and $\mathcal{K}_T(B^c)$ are S -invariant, and therefore S commutes with $e_T(B)$ as well. We prove that, as a consequence of this, $[e_S(A), e_T(B)] = 0$ for every $A \in \mathbb{B}(\mathbb{C})$

3.6 Lemma. *Let $T \in \mathcal{M}$, and let $e \in \mathcal{J}(\tilde{\mathcal{M}})$ with $[e, T] = 0$. Then for every $B \in \mathbb{B}(\mathbb{C})$, $[e, e_T(B)] = 0$. In particular, if $S \in \mathcal{M}$ commutes with T , then $[e_S(\cdot), e_T(\cdot)] = 0$.*

Proof. Let $P = P_{\text{range}(e)}$, $Q = P_{\text{range}(1-e)}$, $R = P_{\text{range}(e_T(B))}$ and $S = P_{\text{range}(1-e_T(B))}$. We prove that (2.26) holds. Since $eT = Te$, $P(\mathcal{H})$ and $Q(\mathcal{H})$ are T -invariant. Then by Lemma 3.3,

$$\begin{aligned} \mathcal{K}_{T|_{P(\mathcal{H})}}(B) &= \mathcal{K}_T(B) \cap P(\mathcal{H}) \\ &= R(\mathcal{H}) \cap P(\mathcal{H}), \end{aligned}$$

and

$$\begin{aligned} \mathcal{K}_{T|_{P(\mathcal{H})}}(B^c) &= \mathcal{K}_T(B^c) \cap P(\mathcal{H}) \\ &= S(\mathcal{H}) \cap P(\mathcal{H}). \end{aligned}$$

Hence $(R \wedge P) \vee (S \wedge P) = P$, and similarly, $(R \wedge Q) \vee (S \wedge Q) = Q$. It follows that

$$1 = P \vee Q = (R \wedge P) \vee (S \wedge P) \vee (R \wedge Q) \vee (S \wedge Q),$$

as desired. $\blacksquare$

Proof of Theorem 3.2. $e_T(\emptyset) = 0$, because $P_T(\emptyset) = 0$. If $B_1, B_2 \in \mathbb{B}(\mathbb{C})$ with $B_1 \cap B_2 = \emptyset$, then $\mathcal{K}_T(B_1) \cap \mathcal{K}_T(B_2) = \{0\}$, i.e. $\text{range}(e_T(B_1)) \cap \text{range}(e_T(B_2)) = \{0\}$. According to Lemma 3.6, $[e_T(B_1), e_T(B_2)] = 0$ so that $e_T(B_1)e_T(B_2) \in \mathcal{J}(\tilde{\mathcal{M}})$. Moreover,

$$\text{range}(e_T(B_1)e_T(B_2)) \subseteq \text{range}(e_T(B_1)) \cap \text{range}(e_T(B_2)) = \{0\},$$

and we conclude that $e_T(B_1)e_T(B_2) = e_T(B_2)e_T(B_1) = 0$.

Now, let $(B_n)_{n=1}^\infty$ be a sequence of mutually disjoint Borel sets. Then for each $N \in \mathbb{N}$ we get from Proposition 3.4 and Lemma 2.7 that

$$\begin{aligned} \text{range}\left(e_T\left(\bigcup_{n=1}^N B_n\right)\right) &= \mathcal{K}_T\left(\bigcup_{n=1}^N B_n\right) \\ &= \overline{\mathcal{K}_T(B_1) + \cdots + \mathcal{K}_T(B_N)} \\ &= \text{range}(e_T(B_1) + \cdots + e_T(B_N)) \end{aligned}$$

and

$$\begin{aligned}
\ker\left(e_T\left(\bigcup_{n=1}^N B_n\right)\right) &= \mathcal{K}_T\left(\left(\bigcup_{n=1}^N B_n\right)^c\right) \\
&= \mathcal{K}_T\left(\bigcap_{n=1}^N B_n^c\right) \\
&= \bigcap_{n=1}^N \mathcal{K}_T(B_n^c) \\
&= \bigcap_{n=1}^N \ker(e_T(B_n)) \\
&= \ker(e_T(B_1) + \cdots + e_T(B_N)).
\end{aligned}$$

Since an element e in $\mathcal{J}(\tilde{\mathcal{M}})$ is uniquely determined by its kernel and its range, it follows that e_T is additive, i.e.

$$e_T\left(\bigcup_{n=1}^N B_n\right) = e_T(B_1) + \cdots + e_T(B_N), \quad (N \in \mathbb{N}). \quad (3.12)$$

Additivity of e_T implies that

$$\begin{aligned}
e_T\left(\bigcup_{n=1}^{\infty} B_n\right) - \sum_{n=1}^{\infty} e_T(B_n) &= \lim_{N \rightarrow \infty} \left(e_T\left(\bigcup_{n=1}^{\infty} B_n\right) - \sum_{n=1}^N e_T(B_n) \right) \\
&= \lim_{N \rightarrow \infty} e_T\left(\bigcup_{n=N+1}^{\infty} B_n\right)
\end{aligned} \quad (3.13)$$

(the limits refer to the measure topology), where

$$\begin{aligned}
\tau\left(\text{supp}\left(e_T\left(\bigcup_{n=N+1}^{\infty} B_n\right)\right)\right) &= \tau\left(P_T\left(\bigcup_{n=N+1}^{\infty} B_n\right)\right) \\
&= \mu_T\left(\bigcup_{n=N+1}^{\infty} B_n\right) \\
&\rightarrow 0, \quad \text{as } N \rightarrow \infty.
\end{aligned}$$

Combining this with (3.13) and Lemma 3.5, we find that e_T is σ -additive as well. $\blacksquare$

Note that in the case where T is a *normal* operator, $B \mapsto P_T(B)$ is just the spectral measure of T , and $e_T(B) = P_T(B)$.

4 The Brown measure of a set of commuting operators in $\mathcal{M}$.

As in the previous section, let $\mathcal{M}$ be a type II_1 factor. The purpose of this section is to prove:

4.1 Theorem. Let $n \in \mathbb{N}$, and let $T_1, \dots, T_n \in \mathcal{M}$ be commuting operators. Then there is a probability measure $\mu_{T_1, \dots, T_n}$ on $(\mathbb{C}^n, \mathbb{B}(\mathbb{C}^n))$, which is uniquely determined by

$$\mu_{T_1, \dots, T_n}(B_1 \times \dots \times B_n) = \tau\left(\bigwedge_{i=1}^n P_{T_i}(B_i)\right), \quad B_1, \dots, B_n \in \mathbb{B}(\mathbb{C}), \quad (4.1)$$

where $P_{T_i}(B_i) \in \mathcal{M}$ is the projection onto $\mathcal{K}_{T_i}(B_i)$ (cf. Section 2).

The idea of proof is as follows:

As mentioned in the previous section (cf. Lemma 3.6), if $S \in \mathcal{M}$ commutes with $T \in \mathcal{M}$, then $[e_S(A), e_T(B)] = 0$ for all $A, B \in \mathbb{B}(\mathbb{C})$. We may therefore define a map $e_{T_1, \dots, T_n}$ from $\mathbb{B}(\mathbb{C})^n$ into $\mathcal{J}(\mathcal{M})$ by

$$e_{T_1, \dots, T_n}(B_1, \dots, B_n) = e_{T_1}(B_1)e_{T_2}(B_2) \cdots e_{T_n}(B_n), \quad B_1, \dots, B_n \in \mathbb{B}(\mathbb{C}). \quad (4.2)$$

We will then define ν on $\mathbb{B}(\mathbb{C})^n$ by

$$\begin{aligned} \nu(B_1, \dots, B_n) &= \tau(\text{supp}[e_{T_1, \dots, T_n}(B_1, \dots, B_n)]) \\ &= \tau(P_{\text{range}(e_{T_1, \dots, T_n}(B_1, \dots, B_n))}) \\ &= \tau\left(\bigwedge_{i=1}^n P_{T_i}(B_i)\right), \quad B_1, \dots, B_n \in \mathbb{B}(\mathbb{C}), \end{aligned} \quad (4.3)$$

and we will prove that ν extends (uniquely) to a probability measure, $\mu_{T_1, \dots, T_n}$, on $(\mathbb{C}^n, \mathbb{B}(\mathbb{C}^n))$.

4.2 Theorem. Consider uncountable, complete, separable metric spaces $(X_1, d_1), \dots, (X_n, d_n)$. Suppose $\nu : \mathbb{B}(X_1) \times \dots \times \mathbb{B}(X_n) \rightarrow [0, \infty[$ is a map satisfying

- (1) for all $B_2 \in \mathbb{B}(X_2), B_3 \in \mathbb{B}(X_3), \dots, B_n \in \mathbb{B}(X_n)$, $B \mapsto \nu(B, B_2, \dots, B_n)$ is a measure on $(X_1, \mathbb{B}(X_1))$,
- (2) for all $B_1 \in \mathbb{B}(X_1), B_3 \in \mathbb{B}(X_3), \dots, B_n \in \mathbb{B}(X_n)$, $B \mapsto \nu(B_1, B, B_3, \dots, B_n)$ is a measure on $(X_2, \mathbb{B}(X_2))$,
- $\vdots$
- (n) for all $B_1 \in \mathbb{B}(X_1), B_2 \in \mathbb{B}(X_2), \dots, B_{n-1} \in \mathbb{B}(X_{n-1})$, $B \mapsto \nu(B_1, B_2, \dots, B_{n-1}, B)$ is a measure on $(X_n, \mathbb{B}(X_n))$.

Then there is a unique measure μ on $\otimes_{i=1}^n \mathbb{B}(X_i)$, such that for all $B_1 \in \mathbb{B}(X_1), B_2 \in \mathbb{B}(X_2), \dots, B_n \in \mathbb{B}(X_n)$,

$$\mu(B_1 \times B_2 \times \dots \times B_n) = \nu(B_1, B_2, \dots, B_n). \quad (4.4)$$

Proof. According to [Ku, Remark 1, p. 358], $(X_i, \mathbb{B}(X_i))$ is Borel equivalent to $([0, 1], \mathbb{B}([0, 1]))$, i.e. there is a bijective bimeasurable map $\phi_i : (X_i, \mathbb{B}(X_i)) \rightarrow ([0, 1], \mathbb{B}([0, 1]))$. Therefore we may as well assume that $X_i = \mathbb{R}$, ($i = 1, \dots, n$). We may also assume that $\nu(\mathbb{R}, \mathbb{R}, \dots, \mathbb{R}) = 1$. We define $F : \mathbb{R}^n \rightarrow [0, 1]$ by

$$F(x_1, \dots, x_n) = \nu([-\infty, x_1], \dots, [-\infty, x_n]), \quad x_1, \dots, x_n \in \mathbb{R}. \quad (4.5)$$

Because of (1)–(n), F is increasing in each variable separately and satisfies

- (a) if $x_i^{(k)} \searrow x_i$, $i = 1, \dots, n$, then $F(x_1^{(k)}, \dots, x_n^{(k)}) \searrow F(x_1, \dots, x_n)$,
- (b) if $x_i \searrow -\infty$ for some $i \in \{1, \dots, n\}$, then $F(x_1, \dots, x_n) \searrow 0$,
- (c) if $x_i \nearrow \infty$ for all $i \in \{1, \dots, n\}$, then $F(x_1, \dots, x_n) \nearrow 1$.

Then, according to [Bre, Corollary 2.27], there is a (unique) probability measure μ on $(\mathbb{R}^n, \mathbb{B}(\mathbb{R}^n))$ such that for all $x_1, \dots, x_n \in \mathbb{R}$,

$$\mu([-\infty, x_1] \times \cdots \times [-\infty, x_n]) = F(x_1, \dots, x_n). \quad (4.6)$$

Let $x_2, \dots, x_n \in \mathbb{R}$ be fixed but arbitrary. Then the (finite) measures

$$B \mapsto \mu(B \times [-\infty, x_2] \times \cdots \times [-\infty, x_n])$$

and

$$B \mapsto \nu(B, [-\infty, x_2], \dots, [-\infty, x_n])$$

have the same distribution functions. Hence they must be identical. That is, for all $B \in \mathbb{B}(\mathbb{R})$,

$$\mu(B \times [-\infty, x_2] \times \cdots \times [-\infty, x_n]) = \nu(B, [-\infty, x_2], \dots, [-\infty, x_n]). \quad (4.7)$$

Now, let $B_1 \in \mathbb{B}(\mathbb{R})$ and $x_3, \dots, x_n \in \mathbb{R}$ be fixed but arbitrary. Then (4.7) shows that the (finite) measures

$$B \mapsto \mu(B_1 \times B \times [-\infty, x_3] \times \cdots \times [-\infty, x_n])$$

and

$$B \mapsto \nu(B_1, B, [-\infty, x_3], \dots, [-\infty, x_n])$$

have the same distribution functions, so they must be identical as well. That is, for all $B \in \mathbb{B}(\mathbb{R})$,

$$\mu(B_1 \times B \times [-\infty, x_3] \times \cdots \times [-\infty, x_n]) = \nu(B_1, B, [-\infty, x_3], \dots, [-\infty, x_n]). \quad (4.8)$$

Continuing like this we find that (4.4) holds. $\blacksquare$

It follows from Theorem 4.2 that in order to show that $\mu_{T_1, \dots, T_n}$ exists, we must prove that (1)–(n) of Theorem 4.2 hold in the case where $X_1 = \cdots = X_n = \mathbb{C}$, and where ν is given by (4.3).

From now on we will, in order to simplify notation a little, consider the case $n = 2$, and we will assume that $S, T \in \mathcal{M}$ are commuting operators. It should be clear that the proof given below may be generalized to the case of arbitrary $n \in \mathbb{N}$.

4.3 Lemma. For fixed $A \in \mathbb{B}(\mathbb{C})$, $\nu_A : \mathbb{B}(\mathbb{C}) \rightarrow [0, 1]$ given by

$$\nu_A(B) = \nu(A, B) = \tau(P_S(A) \wedge P_T(B)), \quad B \in \mathbb{B}(\mathbb{C}) \quad (4.9)$$

(cf. (4.3)) is a measure on $(\mathbb{C}, \mathbb{B}(\mathbb{C}))$.

Proof. According to Theorem 3.2 and Definition 3.1, $e_T(\emptyset) = 0$, so

$$\nu_A(\emptyset) = \tau(\text{supp}[e_S(A)e_T(\emptyset)]) = 0.$$

Let $(B_n)_{n=1}^\infty$ be a sequence of mutually disjoint sets from $\mathbb{B}(\mathbb{C})$. Then $e_T\left(\bigcup_{n=1}^\infty B_n\right) = \sum_{n=1}^\infty e_T(B_n)$, so

$$e_S(A)e_T\left(\bigcup_{n=1}^\infty B_n\right) = \sum_{n=1}^\infty e_S(A)e_T(B_n) \quad (4.10)$$

with $e_S(A)e_T(B_n)e_S(A)e_T(B_m) = e_S(A)e_T(B_n)e_T(B_m) = 0$ when $n \neq m$. Hence, by Proposition 2.8,

$$\mathrm{tr}\left(e_S(A)e_T\left(\bigcup_{n=1}^\infty B_n\right)\right) = \sum_{n=1}^\infty \mathrm{tr}(e_S(A)e_T(B_n)). \quad (4.11)$$

This shows that ν_A is a measure. $\blacksquare$

It now follows from Lemma 4.3 and Theorem 4.2 that there is one and only one (probability) measure $\mu_{S,T}$ on $\mathbb{B}(\mathbb{C}^2)$ such that for all $A, B \in \mathbb{B}(\mathbb{C})$,

$$\mu_{S,T}(A \times B) = \tau(\mathrm{supp}[e_{S,T}(A, B)]) = \tau(P_S(A) \wedge P_T(B)), \quad (4.12)$$

and this proves Theorem 4.1 in the case $n = 2$.

5 Spectral subspaces for commuting operators $S, T \in \mathcal{M}$.

5.1 Theorem. *Let $S, T \in \mathcal{M}$ be commuting operators, and let $B \subseteq \mathbb{C}^2$ be any Borel set. Then there is a maximal, closed, S - and T -invariant subspace $\mathcal{K} = \mathcal{K}_{S,T}(B)$ affiliated with $\mathcal{M}$, such that the Brown measure $\mu_{S|_{\mathcal{K}}, T|_{\mathcal{K}}}$ is concentrated on B . Let $P_{S,T}(B) \in \mathcal{M}$ denote the projection onto $\mathcal{K}_{S,T}(B)$. Then more precisely,*

(i) if $B = B_1 \times B_2$ with $B_1, B_2 \in \mathbb{B}(\mathbb{C})$, then

$$P_{S,T}(B) = P_S(B_1) \wedge P_T(B_2), \quad (5.1)$$

(ii) if B is a disjoint union of the sets $(B_1^{(k)} \times B_2^{(k)})_{k=1}^\infty$, where $B_i^{(k)} \in \mathbb{B}(\mathbb{C})$, $k \in \mathbb{N}$, $i = 1, 2$, then

$$P_{S,T}(B) = \bigvee_{k=1}^\infty [P_S(B_1^{(k)}) \wedge P_T(B_2^{(k)})], \quad (5.2)$$

(iii) and for general $B \in \mathbb{B}(\mathbb{C}^2)$,

$$P_{S,T}(B) = \bigwedge_{B \subseteq U, U \subseteq \mathbb{C}^2 \text{ open}} P_{S,T}(U). \quad (5.3)$$

Moreover,

$$\mu_{S,T}(B) = \tau(P_{S,T}(B)), \quad B \in \mathbb{B}(\mathbb{C}^2). \quad (5.4)$$

5.2 Remark. Every non-empty, open subset of $\mathbb{C}^2 \cong \mathbb{R}^4$ is a disjoint union of countably many *standard intervals*, i.e. sets of the form $\prod_{i=1}^4]a_i, b_i]$, where $-\infty < a_i < b_i < \infty$, $1 \leq i \leq 4$. Hence, it follows from Theorem 5.1 that the map $\mathbb{B}(\mathbb{C}^2) \rightarrow \mathcal{P}(\mathcal{M}) : B \mapsto P_{S,T}(B)$ is uniquely determined by its values on such standard intervals.

Proof of Theorem 5.1. We let $\mathbb{K}$ denote the set of sets of the form $B_1 \times B_2$ with $B_1, B_2 \in \mathbb{B}(\mathbb{C})$.

Consider an arbitrary sequence of mutually disjoint sets from $\mathbb{K}$, $(B_1^{(k)} \times B_2^{(k)})_{k=1}^\infty$, and define

$$P_{S,T} \left(\bigcup_{k=1}^\infty (B_1^{(k)} \times B_2^{(k)}) \right) := \bigvee_{k=1}^\infty [P_S(B_1^{(k)}) \wedge P_T(B_2^{(k)})]. \quad (5.5)$$

Clearly, $P := P_{S,T} \left(\bigcup_{k=1}^\infty (B_1^{(k)} \times B_2^{(k)}) \right)$ satisfies that

(a) P is S - and T -invariant.

In addition, we prove that with $\mathcal{K} = P(\mathcal{H})$,

(b) $\mu_{S|_{\mathcal{K}}, T|_{\mathcal{K}}}$ is concentrated on $B := \bigcup_{k=1}^\infty (B_1^{(k)} \times B_2^{(k)})$,

and

(c) P is maximal w.r.t. the properties (a) and (b).

(c) will entail that the right-hand side of (5.5) is independent of the way in which we write B as a disjoint union of countably many sets from $\mathbb{K}$, and hence, that $P_{S,T}(B)$ *does*, as indicated by the notation, only depend on the set B .

To see that (b) holds, note that if $Q \in \mathcal{M}$ is *any* S - and T -invariant projection, and if we let $\mathcal{L} = Q(\mathcal{H})$, then by (4.1) and Lemma 3.3,

$$\begin{aligned} \mu_{S|_{\mathcal{L}}, T|_{\mathcal{L}}}(B) &= \sum_{k=1}^\infty \mu_{S|_{\mathcal{L}}, T|_{\mathcal{L}}}(B_1^{(k)} \times B_2^{(k)}) \\ &= \sum_{k=1}^\infty \tau_{Q\mathcal{M}Q}(P_{S|_{\mathcal{L}}}(B_1^{(k)}) \wedge P_{T|_{\mathcal{L}}}(B_2^{(k)})) \\ &= \sum_{k=1}^\infty \tau_{Q\mathcal{M}Q}(P_S(B_1^{(k)}) \wedge P_T(B_2^{(k)}) \wedge Q). \end{aligned} \quad (5.6)$$

Then using Proposition 2.8, we get that

$$\begin{aligned} \mu_{S|_{\mathcal{K}}, T|_{\mathcal{K}}}(B) &= \sum_{k=1}^\infty \tau_{P\mathcal{M}P}(P_S(B_1^{(k)}) \wedge P_T(B_2^{(k)}) \wedge P) \\ &= \sum_{k=1}^\infty \tau_{P\mathcal{M}P}(P_S(B_1^{(k)}) \wedge P_T(B_2^{(k)})) \\ &= \frac{1}{\tau(P)} \sum_{k=1}^\infty \text{tr}(e_S(B_1^{(k)})e_T(B_2^{(k)})) \\ &= \frac{1}{\tau(P)} \text{tr} \left(\sum_{k=1}^\infty e_S(B_1^{(k)})e_T(B_2^{(k)}) \right) \\ &= \frac{1}{\tau(P)} \tau \left(\bigvee_{k=1}^\infty [P_S(B_1^{(k)}) \wedge P_T(B_2^{(k)})] \right) \\ &= 1. \end{aligned}$$

Thus, (b) holds.

Now, suppose that $Q \in \mathcal{M}$ is an S - and T -invariant projection, and that $\mu_{S|_{\mathcal{L}}, T|_{\mathcal{L}}}$ is concentrated on B , where $\mathcal{L} = Q(\mathcal{H})$. Then by Lemma 3.3 and Proposition 2.8,

$$\begin{aligned} P \wedge Q &= \left(\bigvee_{k=1}^{\infty} [P_S(B_1^{(k)}) \wedge P_T(B_2^{(k)})] \right) \wedge Q \\ &= \bigvee_{k=1}^{\infty} [P_{S|_{\mathcal{L}}}(B_1^{(k)}) \wedge P_{T|_{\mathcal{L}}}(B_2^{(k)})] \\ &= P_{\text{range}\left(\sum_{k=1}^{\infty} e_{S|_{\mathcal{L}}}(B_1^{(k)}) e_{T|_{\mathcal{L}}}(B_2^{(k)})\right)}. \end{aligned}$$

Hence, Proposition 2.8 and (5.6) imply that

$$\begin{aligned} \tau_{Q\mathcal{M}Q}(P \wedge Q) &= \text{tr}_{Q\mathcal{M}Q}\left(\sum_{k=1}^{\infty} e_{S|_{\mathcal{L}}}(B_1^{(k)}) e_{T|_{\mathcal{L}}}(B_2^{(k)})\right) \\ &= \sum_{k=1}^{\infty} \text{tr}_{Q\mathcal{M}Q}(e_{S|_{\mathcal{L}}}(B_1^{(k)}) e_{T|_{\mathcal{L}}}(B_2^{(k)})) \\ &= \sum_{k=1}^{\infty} \tau_{Q\mathcal{M}Q}(P_{S|_{\mathcal{L}}}(B_1^{(k)}) \wedge P_{T|_{\mathcal{L}}}(B_2^{(k)})) \\ &= \mu_{S|_{\mathcal{L}}, T|_{\mathcal{L}}}(B) \\ &= 1. \end{aligned}$$

Thus, $P \wedge Q = Q$, and this shows that (c) holds.

As mentioned in Remark 5.2, every open set $U \subseteq \mathbb{C}^2$ may be written as a union of countably many mutually disjoint sets from $\mathbb{K}$. Thus, we have now proved existence of $P_{S,T}(U)$ for every such U , and for general $B \in \mathbb{B}(\mathbb{C}^2)$ we will define

$$P_{S,T}(B) := \bigwedge_{B \subseteq U, U \subseteq \mathbb{C}^2 \text{ open}} P_{S,T}(U). \quad (5.7)$$

Then again, $P := P_{S,T}(B)$ satisfies that

(a) P is S - and T -invariant.

Moreover, we prove that with $\mathcal{K} = P(\mathcal{H})$,

(b) $\mu_{S|_{\mathcal{K}}, T|_{\mathcal{K}}}$ is concentrated on B ,

and

(c) P is maximal w.r.t. the properties (a) and (b).

These properties will entail that when B happens to be a union of countably many mutually disjoint sets from $\mathbb{K}$, then (5.7) agrees with the previous definition of $P_{S,T}(B)$ (cf. (5.5)).

Now, to see that (b) holds, note that $\mu_{S|_{\mathcal{K}}, T|_{\mathcal{K}}}$ is regular (cf. [Fo, Theorem 7.8]), and hence

$$\mu_{S|_{\mathcal{K}}, T|_{\mathcal{K}}}(B) = \inf\{\mu_{S|_{\mathcal{K}}, T|_{\mathcal{K}}}(U) \mid B \subseteq U, U \subseteq \mathbb{C}^2 \text{ open}\}. \quad (5.8)$$

Let U be any open subset of $\mathbb{C}^2$ containing B . Write U as a union of countably many mutually disjoint sets from $\mathbb{K}$:

$$U = \bigcup_{k=1}^{\infty} (B_1^{(k)} \times B_2^{(k)}).$$

Then, according to (5.6),

$$\mu_{S|_{\mathcal{K}}, T|_{\mathcal{K}}}(U) = \sum_{k=1}^{\infty} \tau_{PMP}(P_S(B_1^{(k)}) \wedge P_T(B_2^{(k)}) \wedge P),$$

and using Proposition 2.8 and Lemma 3.3 we find that

$$\begin{aligned} \mu_{S|_{\mathcal{K}}, T|_{\mathcal{K}}}(U) &= \text{tr}_{PMP} \left(\sum_{k=1}^{\infty} e_{S|_{\mathcal{K}}}(B_1^{(k)}) e_{T|_{\mathcal{K}}}(B_2^{(k)}) \right) \\ &= \tau_{PMP} \left(\bigvee_{k=1}^{\infty} [P_{S|_{\mathcal{K}}}(B_1^{(k)}) \wedge P_{T|_{\mathcal{K}}}(B_2^{(k)})] \right) \\ &= \tau_{PMP}(P_{S,T}(U) \wedge P) \\ &= \tau_{PMP}(P) \\ &= 1, \end{aligned}$$

where $P_{S,T}(U)$ is given by (5.5). Hence by (5.8), $\mu_{S|_{\mathcal{K}}, T|_{\mathcal{K}}}$ is concentrated on B .

Finally, if $Q \in \mathcal{M}$ is any S - and T -invariant projection, and if $\mu_{S|_{\mathcal{L}}, T|_{\mathcal{L}}}$ is concentrated on B , where $\mathcal{L} = Q(\mathcal{H})$, then $\mu_{S|_{\mathcal{L}}, T|_{\mathcal{L}}}$ is concentrated on U for every open set U containing B . Hence, by the first part of the proof, $Q \leq P_{S,T}(U)$ for every such U , and it follows from the definition of $P_{S,T}(B)$ that $Q \leq P$.

Concerning (5.4), note that if $B = B_1 \times B_2$, where $B_1, B_2 \in \mathbb{B}(\mathbb{C})$, then, by the definitions of $\mu_{S,T}$ and $P_{S,T}(B)$, (5.4) holds. If B is a disjoint union of sets $(B^{(k)})_{k=1}^{\infty} = (B_1^{(k)} \times B_2^{(k)})_{k=1}^{\infty}$, where $B_i^{(k)} \in \mathbb{B}(\mathbb{C})$, $k \in \mathbb{N}$, $i = 1, 2$, then

$$\begin{aligned} \mu_{S,T}(B) &= \sum_{k=1}^{\infty} \tau(P_{S,T}(B_1^{(k)} \times B_2^{(k)})) \\ &= \sum_{k=1}^{\infty} \tau(\text{supp}(e_S(B_1^{(k)}) e_T(B_2^{(k)}))). \end{aligned}$$

Applying Proposition 2.8 we thus find that

$$\begin{aligned} \mu_{S,T}(B) &= \tau \left(\text{supp} \left(\sum_{k=1}^{\infty} e_S(B_1^{(k)}) e_T(B_2^{(k)}) \right) \right) \\ &= \tau \left(\bigvee_{k=1}^{\infty} \text{supp}(e_S(B_1^{(k)}) e_T(B_2^{(k)})) \right) \\ &= \tau \left(\bigvee_{k=1}^{\infty} P_{S,T}(B_1^{(k)} \times B_2^{(k)}) \right) \\ &= \tau(P_{S,T}(B)). \end{aligned}$$

Finally, for general $B \in \mathbb{B}(\mathbb{C}^2)$, since $\mu_{S,T}$ is regular,

$$\begin{aligned}
\mu_{S,T}(B) &= \inf\{\mu_{S,T}(U) \mid B \subseteq U \subseteq \mathbb{C}^2, U \text{ open}\} \\
&= \inf\{\tau(P_{S,T}(U)) \mid B \subseteq U \subseteq \mathbb{C}^2, U \text{ open}\} \\
&= \tau\left(\bigwedge_{B \subseteq U \subseteq \mathbb{C}^2, U \text{ open}} P_{S,T}(U)\right) \\
&= \tau(P_{S,T}(B)). \quad \blacksquare
\end{aligned}$$

The proof given above may clearly be generalized to the case of n commuting operators $T_1, \dots, T_n \in \mathcal{M}$, so that Theorem 5.1 has a slightly more general version:

5.3 Theorem. *Let $n \in \mathbb{N}$, let $T_1, \dots, T_n \in \mathcal{M}$ be commuting operators, and let $B \subseteq \mathbb{C}^n$ be any Borel set. Then there is a maximal closed subspace, $\mathcal{K} = \mathcal{K}_{T_1, \dots, T_n}(B)$, affiliated with $\mathcal{M}$ which is T_i -invariant for every $i \in \{1, \dots, n\}$, and such that the Brown measure $\mu_{T_1|_{\mathcal{K}}, \dots, T_n|_{\mathcal{K}}}$ is concentrated on B . Let $P_{T_1, \dots, T_n}(B) \in \mathcal{M}$ denote the projection onto $\mathcal{K}_{T_1, \dots, T_n}(B)$. Then more precisely,*

(i) *if $B = B_1 \times \dots \times B_n$ with $B_i \in \mathbb{B}(\mathbb{C})$, then*

$$P_{T_1, \dots, T_n}(B) = \bigwedge_{i=1}^n P_{T_i}(B_i), \quad (5.9)$$

(ii) *if B is a disjoint union of sets $(B^{(k)})_{k=1}^\infty = (B_1^{(k)} \times \dots \times B_n^{(k)})_{k=1}^\infty$, where $B_i^{(k)} \in \mathbb{B}(\mathbb{C})$, $k \in \mathbb{N}$, $i = 1, \dots, n$, then*

$$P_{T_1, \dots, T_n}(B) = \bigvee_{k=1}^\infty P_{T_1, \dots, T_n}(B^{(k)}), \quad (5.10)$$

(iii) *and for general $B \in \mathbb{B}(\mathbb{C}^n)$,*

$$P_{T_1, \dots, T_n}(B) = \bigwedge_{B \subseteq U, U \subseteq \mathbb{C}^n \text{ open}} P_{T_1, \dots, T_n}(U). \quad (5.11)$$

Moreover, for every $B \in \mathbb{B}(\mathbb{C}^n)$,

$$\mu_{T_1, \dots, T_n}(B) = \tau(P_{T_1, \dots, T_n}(B)). \quad (5.12)$$

6 An alternative characterization of $\mu_{S,T}$.

In this final section we are going to give a characterization of the Brown measure of two commuting operators in $\mathcal{M}$, which is different from the one we gave in Theorem 4.1. Recall from [Bro] that for $T \in \mathcal{M}$, the Brown measure of T , μ_T , is the unique compactly supported Borel probability measure on $\mathbb{C}$ which satisfies the identity

$$\tau(\log |T - \lambda \mathbf{1}|) = \int_{\mathbb{C}} \log |z - \lambda| d\mu_T(z) \quad (6.1)$$

for all $\lambda \in \mathbb{C}$.

We are going to prove that a similar property characterizes $\mu_{S,T}$:

6.1 Theorem. *Let $S, T \in \mathcal{M}$ be commuting operators. Then $\mu_{S,T}$ is the unique compactly supported Borel probability measure on $\mathbb{C}^2$ which satisfies the identity*

$$\tau(\log |\alpha S + \beta T - \mathbf{1}|) = \int_{\mathbb{C}^2} \log |\alpha z + \beta w - 1| d\mu_{S,T}(z, w). \quad (6.2)$$

for all $\alpha, \beta \in \mathbb{C}$.

6.2 Remark. Let $S, T \in \mathcal{M}$ be as in Theorem 6.1. Note that if $\mu_{S,T}$ satisfies (6.2) for all $\alpha, \beta \in \mathbb{C}$, then for all $\alpha, \beta, \lambda \in \mathbb{C}$,

$$\tau(\log |\alpha S + \beta T - \lambda \mathbf{1}|) = \int_{\mathbb{C}^2} \log |\alpha z + \beta w - \lambda| d\mu_{S,T}(z, w). \quad (6.3)$$

This is clear for $\lambda \neq 0$, and for $\lambda = 0$, (6.3) follows from the fact that two subharmonic functions defined in $\mathbb{C}$ coincide iff they agree almost everywhere w.r.t. Lebesgue measure. It now follows from Brown's characterization of $\mu_{\alpha S + \beta T}$ that $\mu_{\alpha S + \beta T}$ is the push-forward measure $\nu_{\alpha, \beta}$ of $\mu_{S,T}$ via the map $(z, w) \mapsto \alpha z + \beta w$. On the other hand, if $\nu_{\alpha, \beta} = \mu_{\alpha S + \beta T}$, then (6.2) holds.

Recall from [HS] that the *modified spectral radius* of $T \in \mathcal{M}$, $r'(T)$, is defined by

$$r'(T) := \max\{|z| \mid z \in \text{supp}(\mu_T)\}. \quad (6.4)$$

Also recall from [HS, Corollary 2.6] that in fact

$$r'(T) = \lim_{p \rightarrow \infty} \left(\lim_{n \rightarrow \infty} \|T^n\|^{\frac{1}{n}} \right). \quad (6.5)$$

6.3 Lemma. *Let $S, T \in \mathcal{M}$ be commuting operators. Then the modified spectral radii, $r'(S)$, $r'(T)$, $r'(ST)$ and $r'(S + T)$, satisfy the inequalities*

$$r'(ST) \leq r'(S) \cdot r'(T), \quad (6.6)$$

and

$$r'(S + T) \leq r'(S) + r'(T). \quad (6.7)$$

Proof. (6.6) follows from (6.5) and the generalized Hölder inequality (cf. [FK]): For $A, B \in \mathcal{M}$ and for $0 < p, q, r \leq \infty$ with $\frac{1}{r} = \frac{1}{p} + \frac{1}{q}$,

$$\|AB\|_r \leq \|A\|_p \|B\|_q.$$

To prove (6.7), note that

$$\text{supp}(\mu_S) \subseteq \{z \in \mathbb{C} \mid \text{Re } z \leq r'(S)\},$$

and

$$\text{supp}(\mu_T) \subseteq \{z \in \mathbb{C} \mid \text{Re } z \leq r'(T)\}.$$

According to [Bro, Theorem 4.1], μ_{e^S} and μ_{e^T} are the push-forward measures of μ_S and μ_T , respectively, via the map $z \mapsto e^z$. Hence,

$$\text{supp}(\mu_{e^S}) \subseteq \{z \in \mathbb{C} \mid |z| \leq e^{r'(S)}\},$$

and

$$\text{supp}(\mu_{e^T}) \subseteq \{z \in \mathbb{C} \mid |z| \leq e^{r'(T)}\},$$

and it follows from (6.6) that

$$\begin{aligned} r'(e^{S+T}) &= r'(e^S e^T) \\ &\leq e^{r'(S)} e^{r'(T)} \\ &= e^{r'(S)+r'(T)}. \end{aligned}$$

Thus, $\text{supp}(\mu_{e^{S+T}}) \subseteq \overline{B(0, e^{r'(S)+r'(T)})}$, and then, by one more application of [Bro, Theorem 4.1],

$$\text{supp}(\mu_{S+T}) \subseteq \{z \in \mathbb{C} \mid \text{Re} z \leq r'(S) + r'(T)\}.$$

Repeating this argument, we find that for arbitrary $\theta \in [0, 2\pi[$,

$$\text{supp}(\mu_{e^{i\theta}(S+T)}) \subseteq \{z \in \mathbb{C} \mid \text{Re} z \leq r'(S) + r'(T)\},$$

i.e.

$$\text{supp}(\mu_{S+T}) \subseteq \{z \in \mathbb{C} \mid \text{Re}(e^{-i\theta} z) \leq r'(S) + r'(T)\}.$$

Since θ was arbitrary, we conclude that

$$\text{supp}(\mu_{S+T}) \subseteq \overline{B(0, r'(S) + r'(T))},$$

and this proves (6.7). $\blacksquare$

6.4 Lemma. Let $S, T \in \mathcal{M}$ be commuting operators, and let $\alpha, \beta \in \mathbb{C}$. Then $\mu_{\alpha S, \beta T}$ is the push-forward measure of $\mu_{S, T}$ via the map $h_{\alpha, \beta} : \mathbb{C} \times \mathbb{C} \rightarrow \mathbb{C} \times \mathbb{C}$ given by

$$h_{\alpha, \beta}(z, w) = (\alpha z, \beta w).$$

Proof. Recall that $\mu_{\alpha S, \beta T}$ is uniquely determined by the property that for all $B_1, B_2 \in \mathbb{B}(\mathbb{C})$,

$$\mu_{\alpha S, \beta T}(B_1 \times B_2) = \tau(P_{\alpha S}(B_1) \wedge P_{\beta T}(B_2)). \quad (6.8)$$

Now, it is easily seen that for $\alpha \neq 0$ and $\beta \neq 0$, $P_{\alpha S}(B_1) = P_S(\frac{1}{\alpha} B_1)$ and $P_{\beta T}(B_2) = P_T(\frac{1}{\beta} B_2)$. Hence,

$$\begin{aligned} \mu_{\alpha S, \beta T}(B_1 \times B_2) &= \tau(P_S(\frac{1}{\alpha} B_1) \wedge P_T(\frac{1}{\beta} B_2)) \\ &= \mu_{S, T}(\frac{1}{\alpha} B_1 \times \frac{1}{\beta} B_2) \\ &= \mu_{S, T}(h_{\alpha, \beta}^{-1}(B_1 \times B_2)). \end{aligned} \quad (6.9)$$

If for instance $\alpha = 0$, then $P_{\alpha S}(B_1) = 0$ if $0 \notin B_1$ and $P_{\alpha S}(B_1) = \mathbf{1}$ if $0 \in B_1$. It then follows that (6.9) holds in this case as well. Similar arguments apply if $\beta = 0$. $\blacksquare$

Proof of Theorem 6.1. As noted in Remark 6.2, it suffices to prove that for all $\alpha, \beta \in \mathbb{C}$, $\mu_{\alpha S + \beta T}$ is the push-forward measure of $\mu_{S, T}$ via the map $(z, w) \mapsto \alpha z + \beta w$. At first we will consider the case $\alpha = \beta = 1$. Define $a : \mathbb{C} \times \mathbb{C} \rightarrow \mathbb{C}$ by

$$a(z, w) = z + w, \quad (z, w \in \mathbb{C}).$$

We are going to prove that for all $B \in \mathbb{B}(\mathbb{C})$,

$$\mu_{S+T}(B) = \mu_{S,T}(a^{-1}(B)). \quad (6.10)$$

It suffices to show that for every open set $U \subseteq \mathbb{C}$,

$$\mu_{S+T}(U) \geq \mu_{S,T}(a^{-1}(U)). \quad (6.11)$$

Indeed, if this holds, then by regularity of μ_{S+T} and $a(\mu_{S,T})$, for every Borel set $B \subseteq \mathbb{C}$,

$$\begin{aligned} \mu_{S+T}(B) &= \inf\{\mu_{S+T}(U) \mid B \subseteq U, U \text{ open}\} \\ &\geq \inf\{a(\mu_{S,T})(U) \mid B \subseteq U, U \text{ open}\} \\ &= \mu_{S,T}(a^{-1}B). \end{aligned}$$

Since both measures are probability measures, and the above inequality holds for both B and B^c , we must have identity. That is, (6.10) holds.

Now, let $U \subseteq \mathbb{C}$ be any open set. Then $V := a^{-1}(U)$ is open in $\mathbb{C}^2$ and we may write V as a countable union of mutually disjoint "boxes",

$$V = \bigcup_{n=1}^{\infty} I(z_n, \delta_n) \times I(w_n, \delta_n),$$

where for $z \in \mathbb{C}$ and $\delta > 0$,

$$I(z, \delta) := \{w \in \mathbb{C} \mid \operatorname{Re}(z) - \delta < \operatorname{Re}(w) \leq \operatorname{Re}(z) + \delta, \operatorname{Im}(z) - \delta < \operatorname{Im}(w) \leq \operatorname{Im}(z) + \delta\}. \quad (6.12)$$

We can even choose z_n, w_n and $\delta_n > 0$ so that

$$\overline{B(z_n + w_n, 2\sqrt{2}\delta_n)} \subseteq U. \quad (6.13)$$

This requires a little consideration and for the convenience of the reader, we provide an argument in Lemma 6.7 below.

Now, according to Theorem 5.1,

$$\begin{aligned} \mu_{S,T}(a^{-1}(U)) &= \tau(P_{S,T}(V)) \\ &= \tau\left(\bigvee_{n=1}^{\infty} [P_S(I(z_n, \delta_n)) \wedge P_T(I(w_n, \delta_n))]\right), \end{aligned}$$

and we also have that

$$\mu_{S+T}(U) = \tau(P_{S+T}(U)).$$

Hence, it suffices to prove that for every $n \in \mathbb{N}$,

$$P_{S+T}(U) \geq P_S(I(z_n, \delta_n)) \wedge P_T(I(w_n, \delta_n)). \quad (6.14)$$

Fix $n \in \mathbb{N}$, and set $P = P_S(I(z_n, \delta_n)) \wedge P_T(I(w_n, \delta_n))$. Then by Lemma 6.3,

$$\begin{aligned} r'([S + T - (z_n + w_n)\mathbf{1}]|_{P(\mathcal{H})}) &\leq r'([S - z_n\mathbf{1}]|_{P(\mathcal{H})}) + r'([T - w_n\mathbf{1}]|_{P(\mathcal{H})}) \\ &\leq r'([S - z_n\mathbf{1}]|_{P_S(I(z_n, \delta_n))(\mathcal{H})}) + r'([T - w_n\mathbf{1}]|_{P_T(I(w_n, \delta_n))(\mathcal{H})}) \\ &\leq 2\sqrt{2}\delta_n, \end{aligned}$$

and it follows that $\mu_{S+T|P(\mathfrak{H})}$ is concentrated on $\overline{B(z_n + w_n, 2\sqrt{2}\delta_n)} \subseteq U$. Hence, $P \leq P_{S+T}(U)$, and we are done.

Now, if $\alpha, \beta \in \mathbb{C}$, then we conclude from the above and Lemma 6.4 that

$$\begin{aligned}\mu_{\alpha S + \beta T}(B) &= \mu_{\alpha S, \beta T}(a^{-1}(B)) \\ &= \mu_{S, T}(h_{\alpha, \beta}^{-1}(a^{-1}(B))) \\ &= \mu_{S, T}((a \circ h_{\alpha, \beta})^{-1}(B)),\end{aligned}$$

and since $(a \circ h_{\alpha, \beta})(z, w) = \alpha z + \beta w$, this completes the proof of the identity (6.2).

To prove uniqueness of $\mu_{S, T}$, suppose that ν is a compactly supported Borel probability measure on $\mathbb{C}^2$ which satisfies the identity (6.2) for all $\alpha, \beta \in \mathbb{C}$. That is, for all $\alpha, \beta \in \mathbb{C}$, $\mu_{\alpha S + \beta T}$ is the push-forward measure of ν via the map $(z, w) \mapsto \alpha z + \beta w$. Then, to prove that $\nu = \mu_{S, T}$, it suffices to prove that for all $y = (y_1, \dots, y_4) \in \mathbb{R}^4$,

$$\int_{\mathbb{R}^4} e^{i(y, x)} d\mu_{S, T}(x) = \int_{\mathbb{R}^4} e^{i(y, x)} d\nu(x) \quad (6.15)$$

(here we identify $\mathbb{C}$ with $\mathbb{R}^2$). For $x = (x_1, \dots, x_4) \in \mathbb{R}^4$ and $y = (y_1, \dots, y_4) \in \mathbb{R}^4$, note that

$$(y, x) = \operatorname{Re}\left((y_1 - iy_2)(x_1 + ix_2) + (y_3 - iy_4)(x_3 + ix_4)\right),$$

and hence with $\alpha = y_1 - iy_2$ and $\beta = y_3 - iy_4$ we find that

$$\begin{aligned}\int_{\mathbb{R}^4} e^{i(y, x)} d\mu_{S, T}(x) &= \int_{\mathbb{C}^2} e^{i\operatorname{Re}(\alpha z + \beta w)} d\mu_{S, T}(z, w) \\ &= \int_{\mathbb{C}} e^{i\operatorname{Re} z} d\mu_{\alpha S + \beta T}(z) \\ &= \int_{\mathbb{C}^2} e^{i\operatorname{Re}(\alpha z + \beta w)} d\nu(z, w) \\ &= \int_{\mathbb{R}^4} e^{i(y, x)} d\nu(x),\end{aligned}$$

as desired. $\blacksquare$

6.5 Remark. In the proof above it was shown that for $U \subseteq \mathbb{C}$ an open set, we have the following inequality:

$$P_{S+T}(U) \geq P_{S, T}(a^{-1}(U)). \quad (6.16)$$

But it was also shown that the two projections above have the same trace:

$$\tau(P_{S+T}(U)) = \mu_{S+T}(U) = \mu_{S, T}(a^{-1}(U)) = \tau(P_{S, T}(a^{-1}(U))).$$

Hence, the two projections in (6.16) are identical, and by Theorem 5.1 (iii), for *every* Borel set $B \subseteq \mathbb{C}$, we must have that

$$P_{S+T}(B) = P_{S, T}(p^{-1}(B)). \quad (6.17)$$

As in the previous section, one can easily generalize the proof given above to the case of an arbitrary finite set of commuting operators, $\{T_1, \dots, T_n\}$. That is, we actually have the following alternative description of $\mu_{T_1, \dots, T_n}$:

6.6 Theorem. Let $n \in \mathbb{N}$, and let $T_1, \dots, T_n$ be mutually commuting operators in $\mathcal{M}$. Then $\mu_{T_1, \dots, T_n}$ is the unique compactly supported Borel probability measure on $\mathbb{C}^n$ which satisfies the identity

$$\tau(\log |\alpha_1 T_1 + \dots + \alpha_n T_n - \mathbf{1}|) = \int_{\mathbb{C}^n} \log |\alpha_1 z_1 + \dots + \alpha_n z_n - 1| d\mu_{T_1, \dots, T_n}(z_1, \dots, z_n) \quad (6.18)$$

for all $\alpha_1, \dots, \alpha_n \in \mathbb{C}$.

6.7 Lemma. Define $a : \mathbb{C}^2 \rightarrow \mathbb{C}$ by

$$a(z, w) = z + w,$$

and let $U \subseteq \mathbb{C}$ be an open set. Then for every pair (S, T) of commuting operators in $\mathcal{M}$, we may write $V := a^{-1}(U)$ as a countable disjoint union of sets $(I(z_n, \delta_n) \times I(w_n, \delta_n))_{n=1}^\infty$, where for $z \in \mathbb{C}$ and $\delta > 0$,

$$I(z, \delta) := \{w \in \mathbb{C} \mid \operatorname{Re}(z) - \delta < \operatorname{Re}(w) \leq \operatorname{Re}(z) + \delta, \operatorname{Im}(z) - \delta < \operatorname{Im}(w) \leq \operatorname{Im}(z) + \delta\}.$$

Moreover, we can ensure that for each $n \in \mathbb{N}$,

$$\overline{B(z_n + w_n, 2\sqrt{2}\delta_n)} \subseteq U. \quad (6.19)$$

Proof. Divide $\mathbb{C}^2$ into mutually disjoint "boxes" of the form $I(z, 1) \times I(w, 1)$. Take the countably many of these, $(I(z_n^{(1)}, 1) \times I(w_n^{(1)}, 1))$, which are contained in V and satisfy (6.19) (with $\delta_n = 1$). Next, divide $\mathbb{C}^2$ into boxes once more by cutting each of the previous ones into 16 boxes of equal size (edge length $\frac{1}{2}$). Take those, $(I(z_n^{(2)}, \frac{1}{2}) \times I(w_n^{(2)}, \frac{1}{2}))$, which are not contained in $\bigcup_n I(z_n^{(1)}, 1) \times I(w_n^{(1)}, 1)$ but which are contained in V and satisfy (6.19) (with $\delta_n = \frac{1}{2}$). Continue like this and obtain a set $V_0 \subseteq V$ of the form

$$V_0 = \bigcup_{m=1}^\infty \bigcup_n I(z_n^{(m)}, 2^{-m}) \times I(w_n^{(m)}, 2^{-m}).$$

We claim that this disjoint union is all of V . Indeed, let $(z, w) \in V$. Then $z + w \in U$ so there is an $\varepsilon > 0$ such that $\overline{B(z + w, \varepsilon)} \subseteq U$. Choose $m \in \mathbb{N}$ so large that $2^{-m} < \frac{\varepsilon}{4\sqrt{2}}$. Then, if $|z - z'| \leq \sqrt{2}2^{-m}$ and $|w - w'| \leq \sqrt{2}2^{-m}$, one has that

$$\overline{B(z' + w', 2\sqrt{2}2^{-m})} \subseteq \overline{B(z + w, 4\sqrt{2}2^{-m})} \subseteq U.$$

Thus, when we have divided $\mathbb{C}^2$ into boxes of edge length at most 2^{-m} , then the one box containing (z, w) will satisfy (6.19). Then it is just a matter of taking m so large that the box is also contained in V . It follows that $(z, w) \in V_0$. ■

6.8 Remark. Let $S \in \mathcal{M}$ be invertible, and let $T_1, \dots, T_n$ be mutually commuting operators in $\mathcal{M}$. Then

$$\mu_{S^{-1}T_1S, \dots, S^{-1}T_nS} = \mu_{T_1, \dots, T_n}. \quad (6.20)$$

Indeed, this follows from the characterization of $\mu_{T_1, \dots, T_n}$ given in Theorem 6.6 and from the fact that for all $T \in \mathcal{M}$, $\mu_{S^{-1}TS} = \mu_T$ (cf. [Bro]).

6.9 Proposition. *Let $S, T \in \mathcal{M}$ be commuting operators. Then μ_{ST} is the push-forward measure of $\mu_{S,T}$ via the map $m : (z, w) \mapsto zw$.*

Proof. The proof is essentially the same as the one we gave above when considering the map $a : (z, w) \mapsto z + w$. Again it suffices to show that for every open set $U \subseteq \mathbb{C}$,

$$\mu_{ST}(U) \geq \mu_{S,T}(m^{-1}(U)), \quad (6.21)$$

and for such an open set U we write $V := m^{-1}(U)$ as a countable union of mutually disjoint "boxes" as in (6.12), but this time we make sure that $\delta_n > 0$ is so small that

$$\overline{B(z_n w_n, \sqrt{2}\delta_n(\|T\| + |z_n|))} \subseteq U. \quad (6.22)$$

As in the previous case, one only has to show that for every $n \in \mathbb{N}$,

$$P_{ST}(U) \geq P_S(I(z_n, \delta_n)) \wedge P_T(I(w_n, \delta_n)).$$

Fix $n \in \mathbb{N}$ and set $P = P_S(I(z_n, \delta_n)) \wedge P_T(I(w_n, \delta_n))$. Since

$$r'([S - z_n \mathbf{1}]|_{P(\mathcal{H})}) \leq r'([S - z_n \mathbf{1}]|_{P_S(I(z_n, \delta_n))(\mathcal{H})}) \leq \sqrt{2}\delta_n,$$

and

$$r'([T - w_n \mathbf{1}]|_{P(\mathcal{H})}) \leq r'([T - w_n \mathbf{1}]|_{P_T(I(w_n, \delta_n))(\mathcal{H})}) \leq \sqrt{2}\delta_n,$$

and since

$$ST - z_n w_n \mathbf{1} = (S - z_n \mathbf{1})T + z_n(T - w_n \mathbf{1}),$$

we have (cf. Lemma 6.3) that

$$\begin{aligned} r'([ST - z_n w_n \mathbf{1}]|_{P(\mathcal{H})}) &\leq r'([(S - z_n \mathbf{1})T]|_{P(\mathcal{H})}) + |z_n| r'([T - w_n \mathbf{1}]|_{P(\mathcal{H})}) \\ &\leq r'([S - z_n \mathbf{1}]|_{P(\mathcal{H})})\|T\| + |z_n| r'([T - w_n \mathbf{1}]|_{P(\mathcal{H})}) \\ &\leq \sqrt{2}\delta_n(\|T\| + |z_n|). \end{aligned}$$

Thus, $\mu_{ST|_{P(\mathcal{H})}}$ is concentrated on $\overline{B(z_n w_n, \sqrt{2}\delta_n(\|T\| + |z_n|))} \subseteq U$, and therefore $P \leq P_{ST}(U)$, as desired. ■

6.10 Remark. As in the additive case, we infer from the proof given above that for every Borel set $B \subseteq \mathbb{C}$ we have:

$$P_{ST}(B) = P_{S,T}(m^{-1}(B)). \quad (6.23)$$

6.11 Proposition. *Consider type II_1 factors $\mathcal{M}_1$ and $\mathcal{M}_2$ with faithful tracial states τ_1 and τ_2 , respectively. Let $S \in \mathcal{M}_1$ and $T \in \mathcal{M}_2$. Then*

$$\mu_{S \otimes \mathbf{1}, \mathbf{1} \otimes T} = \mu_S \otimes \mu_T, \quad (6.24)$$

and it follows that

$$\mu_{S \otimes \mathbf{1} + \mathbf{1} \otimes T} = \mu_S * \mu_T, \quad (6.25)$$

$$\mu_{S \otimes T} = \mu_S \star \mu_T, \quad (6.26)$$

where $*$ ($\star$, resp.) denotes additive (multiplicative, resp.) convolution.

Proof. $\mu_{S \otimes \mathbf{1} + \mathbf{1} \otimes T}$ ($\mu_{S \otimes T}$, resp.) is the push-forward measure of $\mu_{S \otimes \mathbf{1}, \mathbf{1} \otimes T}$ via the map $a : (z, w) \mapsto z + w$ ($m : (z, w) \mapsto zw$, resp.), and $\mu_S * \mu_T$ ($\mu_S \star \mu_T$, resp.) is the push-forward measure of $\mu_S \otimes \mu_T$ via that same map. Thus, (6.25) and (6.26) follow from (6.24). To see that the latter holds, let $B_1, B_2 \in \mathbb{B}(\mathbb{C})$. It is easily seen that

$$P_{S \otimes \mathbf{1}}(B_1) = P_S(B_1) \otimes \mathbf{1} \quad \text{and} \quad P_{\mathbf{1} \otimes T}(B_2) = \mathbf{1} \otimes P_T(B_2).$$

Hence,

$$\begin{aligned} \mu_{S \otimes \mathbf{1}, \mathbf{1} \otimes T}(B_1 \times B_2) &= (\tau_1 \otimes \tau_2)[(P_S(B_1) \otimes \mathbf{1}) \cap (\mathbf{1} \otimes P_T(B_2))] \\ &= \tau_1(P_S(B_1))\tau_2(P_T(B_2)) \\ &= \mu_S(B_1)\mu_T(B_2) \\ &= (\mu_S \otimes \mu_T)(B_1 \times B_2). \end{aligned}$$

This proves (6.24). $\blacksquare$

7 Polynomials in n commuting variables.

In this final section we will prove:

7.1 Theorem. *Let $n \in \mathbb{N}$, and let q be a polynomial in n commuting variables, i.e. $q \in \mathbb{C}[z_1, \dots, z_n]$. Then for every n -tuple $(T_1, \dots, T_n)$ of commuting operators in $\mathcal{M}$, one has that*

$$\mu_{q(T_1, \dots, T_n)} = q(\mu_{T_1, \dots, T_n}), \quad (7.1)$$

where $q(\mu_{T_1, \dots, T_n})$ is the push-forward measure of $\mu_{T_1, \dots, T_n}$ via $q : \mathbb{C}^n \rightarrow \mathbb{C}$.

The proof relies on the previous sections and a few technical lemmas.

7.2 Lemma. *Given $n \in \mathbb{N}$ and commuting operators $T_1, \dots, T_n \in \mathcal{M}$. Let $1 \leq i < n$. Then for all Borel sets $A \subseteq \mathbb{C}^i$ and $B \subseteq \mathbb{C}^{n-i}$, one has that*

$$P_{T_1, \dots, T_n}(A \times B) = P_{T_1, \dots, T_i}(A) \wedge P_{T_{i+1}, \dots, T_n}(B). \quad (7.2)$$

Proof. Fix Borel sets $A_1, \dots, A_i \subseteq \mathbb{C}$, put $A = A_1 \times \dots \times A_i$, and let $B \subseteq \mathbb{C}^{n-i}$ be any open set. Then we may write B as a disjoint union of cartesian products of Borel sets $B_m^{(j)}$, $i+1 \leq j \leq n$, i.e.

$$B = \bigcup_{m=1}^{\infty} [B_m^{(i+1)} \times \dots \times B_m^{(n)}].$$

Then by Theorem 5.3, (7.2) holds. Now, let $B \subseteq \mathbb{C}^{n-i}$ be any Borel set. Then by Theorem 5.3,

$$P_{T_{i+1}, \dots, T_n}(B) = \bigwedge_{B \subseteq U, U \text{ open}} P_{T_{i+1}, \dots, T_n}(U),$$

and it follows from the above that

$$P_{T_1, \dots, T_i}(A) \wedge P_{T_{i+1}, \dots, T_n}(B) = \bigwedge_{B \subseteq U, U \text{ open}} P_{T_1, \dots, T_i}(A) \wedge P_{T_{i+1}, \dots, T_n}(U) \quad (7.3)$$

$$= \bigwedge_{B \subseteq U, U \text{ open}} P_{T_1, \dots, T_n}(A \times U). \quad (7.4)$$

Since $P_{T_1, \dots, T_n}(A \times B) \leq P_{T_1, \dots, T_n}(A \times U)$ when $B \subseteq U$, this implies that

$$P_{T_1, \dots, T_i}(A) \wedge P_{T_{i+1}, \dots, T_n}(B) \geq P_{T_1, \dots, T_n}(A \times B). \quad (7.5)$$

On the other hand, (7.4) shows that

$$\begin{aligned} \tau(P_{T_1, \dots, T_i}(A) \wedge P_{T_{i+1}, \dots, T_n}(B)) &= \inf_{B \subseteq U, U \text{ open}} \tau(P_{T_1, \dots, T_n}(A \times U)) \\ &= \inf_{B \subseteq U, U \text{ open}} \mu_{T_1, \dots, T_n}(A \times U). \end{aligned}$$

For fixed A as above, the map $B \mapsto \mu_{T_1, \dots, T_n}(A \times B)$ defined on Borel subsets of $\mathbb{C}^{n-i}$ is a finite (hence regular) Borel measure. Therefore we conclude that

$$\begin{aligned} \tau(P_{T_1, \dots, T_i}(A) \wedge P_{T_{i+1}, \dots, T_n}(B)) &= \mu_{T_1, \dots, T_n}(A \times B) \\ &= \tau(P_{T_1, \dots, T_n}(A \times B)). \end{aligned}$$

Comparing this identity with (7.5), we find that (7.2) holds for this particular A . Now fix an arbitrary Borel set $B \subseteq \mathbb{C}^{n-i}$, and proceed in the same manner. That is, at first assume that $A \subseteq \mathbb{C}^i$ is open and verify that (7.2) holds in this case. Then finally consider an arbitrary Borel set A . ■

7.3 Lemma. Let $n \in \mathbb{N}$ and let $\alpha \in \mathbb{C}$. Define $a_n, m_n^{(\alpha)} : \mathbb{C}^{n+1} \rightarrow \mathbb{C}$ by

$$a_n(z_1, \dots, z_n, z_{n+1}) = (z_1, \dots, z_n + z_{n+1}), \quad (7.6)$$

$$m_n^{(\alpha)}(z_1, \dots, z_n, z_{n+1}) = (z_1, \dots, \alpha z_n z_{n+1}). \quad (7.7)$$

Then for any $(n+1)$ -tuple $(T_1, \dots, T_{n+1})$ of commuting operators in $\mathcal{M}$ one has that

$$\mu_{T_1, \dots, T_{n-1}, T_n + T_{n+1}} = a_n(\mu_{T_1, \dots, T_{n+1}}), \quad (7.8)$$

and

$$\mu_{T_1, \dots, T_{n-1}, \alpha T_n T_{n+1}} = m_n^{(\alpha)}(\mu_{T_1, \dots, T_{n+1}}). \quad (7.9)$$

Proof. The proof is based on Lemma 7.2 and the fact that by (6.17) and (6.23), for any Borel set $B \subseteq \mathbb{C}$ we have:

$$P_{T_n + T_{n+1}}(B) = P_{T_n, T_{n+1}}(1^{-1}(B)) \quad (7.10)$$

and

$$P_{\alpha T_n T_{n+1}}(B) = P_{\alpha T_n, T_{n+1}}(m^{-1}(B)) = P_{T_n, T_{n+1}}((m^{(\alpha)})^{-1}(B)).$$

In order to prove (7.8), consider arbitrary Borel sets $B_1, \dots, B_n \subseteq \mathbb{C}$. We must show that

$$\mu_{T_1, \dots, T_{n-1}, T_n + T_{n+1}}(B_1 \times \dots \times B_n) = \mu_{T_1, \dots, T_{n+1}}(a_n^{-1}(B_1 \times \dots \times B_n)),$$

i.e. that

$$\mu_{T_1, \dots, T_{n-1}, T_n + T_{n+1}}(B_1 \times \dots \times B_n) = \mu_{T_1, \dots, T_{n+1}}(B_1 \times \dots \times B_{n-1} \times a^{-1}(B_n)). \quad (7.11)$$

But by Lemma 7.2 and by (7.10),

$$\begin{aligned} P_{T_1, \dots, T_{n-1}, T_n + T_{n+1}}(B_1 \times \dots \times B_n) &= P_{T_1}(B_1) \wedge \dots \wedge P_{T_{n-1}}(B_{n-1}) \wedge P_{T_n + T_{n+1}}(B_n) \\ &= P_{T_1}(B_1) \wedge \dots \wedge P_{T_{n-1}}(B_{n-1}) \wedge P_{T_n, T_{n+1}}(a^{-1}(B_n)) \\ &= P_{T_1, \dots, T_{n+1}}(B_1 \times \dots \times B_{n-1} \times a^{-1}(B_n)), \end{aligned}$$

and this proves (7.11). (7.9) follows in a similar way. ■

7.4 Lemma. Let $n \in \mathbb{N}$, and let $\sigma \in S_n$ (the group of permutations of $\{1, 2, \dots, n\}$). Then for any n -tuple $(T_1, \dots, T_n)$ of commuting operators in $\mathcal{M}$,

$$\mu_{T_{\sigma(1)}, \dots, T_{\sigma(n)}} = \sigma(\mu_{T_1, \dots, T_n}), \quad (7.12)$$

where identify σ with the corresponding permutation of coordinates $\mathbb{C}^n \rightarrow \mathbb{C}^n$.

Proof. This follows easily from Theorem 4.1. $\blacksquare$

7.5 Lemma. For $n \in \mathbb{N}$ and $1 \leq i \leq n$ define $f_i : \mathbb{C}^n \rightarrow \mathbb{C}^{n+1}$ by

$$f_i(z_1, \dots, z_n) = (z_1, \dots, z_n, z_i).$$

Then for n commuting operators $T_1, \dots, T_n \in \mathcal{M}$, one has that

$$\mu_{T_1, \dots, T_n, T_i} = f_i(\mu_{T_1, \dots, T_n}). \quad (7.13)$$

Proof. Given Borel sets $B_1, \dots, B_{n+1} \subseteq \mathbb{C}$ we must show that

$$\mu_{T_1, \dots, T_n, T_i}(B_1 \times \dots \times B_{n+1}) = \mu_{T_1, \dots, T_n}(f_i^{-1}(B_1 \times \dots \times B_{n+1})). \quad (7.14)$$

Clearly,

$$f_i^{-1}(B_1 \times \dots \times B_{n+1}) = B_1 \times \dots \times (B_i \cap B_{n+1}) \times \dots \times B_n$$

so that the right-hand side of (7.14) is

$$\begin{aligned} \tau(P_{T_1}(B_1) \wedge \dots \wedge P_{T_i}(B_i \cap B_{n+1}) \wedge \dots \wedge P_{T_n}(B_n)) = \\ \tau(P_{T_1}(B_1) \wedge \dots \wedge P_{T_i}(B_i) \wedge P_{T_i}(B_{n+1}) \wedge \dots \wedge P_{T_n}(B_n)). \end{aligned}$$

But this is exactly the left-hand side of (7.14) and we are done. $\blacksquare$

We will not give the proof of Theorem 7.1 in full generality but rather, by way of an example, illustrate how it goes. Consider for instance 3 commuting operators $T_1, T_2, T_3 \in \mathcal{M}$ and the polynomial $q \in \mathbb{C}[z_1, z_2, z_3]$ given by

$$q(z_1, z_2, z_3) = 1 + 2z_2^2 + z_1 z_2 z_3. \quad (7.15)$$

At first define $\phi_1 : \mathbb{C}^3 \rightarrow \mathbb{C}^5$ by

$$\phi_1(z_1, z_2, z_3) = (z_2, z_2, z_1, z_2, z_3).$$

By repeated use of Lemma 7.4 and Lemma 7.5 we find that

$$\mu_{T_2, T_2, T_1, T_2, T_3} = \phi_1(\mu_{T_1, T_2, T_3}).$$

Next define $\phi_2 : \mathbb{C}^5 \rightarrow \mathbb{C}^2$ by

$$\phi_2(z_1, \dots, z_5) = (2z_1 z_2, z_3 z_4 z_5),$$

and by repeated use of (7.9) and Lemma 7.4 conclude that

$$\mu_{2T_2^2, T_1 T_2 T_3} = (\phi_2 \circ \phi_1)(\mu_{T_1, T_2, T_3}).$$

With $\phi_3 : \mathbb{C}^2 \rightarrow \mathbb{C}$ given by

$$\phi_3(z_1, z_2) = z_1 + z_2$$

we now have (cf. (7.8)) that

$$\mu_{(q-1)(T_1, T_2, T_3)} = (\phi_3 \circ \phi_2 \circ \phi_1)(\mu_{T_1, T_2, T_3}) = (q-1)(\mu_{T_1, T_2, T_3}).$$

It is now a simple matter to show that for all $\lambda \in \mathbb{C}$,

$$\tau(\log |q(T_1, T_2, T_3) - \lambda \mathbf{1}|) = \int_{\mathbb{C}} \log |z - \lambda| dq(\mu_{T_1, T_2, T_3})(z),$$

and then by Brown's characterization of $\mu_{q(T_1, T_2, T_3)}$, $\mu_{q(T_1, T_2, T_3)} = q(\mu_{T_1, T_2, T_3})$, as desired.

References

- [A] P. Ainsworth, Ubegrænsede operatorer affilieret med en endelig von Neumann algebra, Master Thesis, University of Odense, 1985.
- [Aa] L. Aagaard, The non-microstates free entropy dimension of DT-operators, *Journal of Functional Analysis* (2004), –.
- [Bre] L. Breiman, Probability, *Addison-Wesley Publishing Company*, 1968.
- [Bro] L. G. Brown, Lidskii's Theorem in the Type II Case, Geometric methods in operator algebras (Kyoto 1983), H. Araki and E. Effros (Eds.) *Pitman Res. notes in Math.* Ser 123, Longman Sci. Tech. (1986), 1–35.
- [FK] T. Fack, H. Kosaki, Generalized s -numbers of τ -measurable operators, *Pacific Journal of Mathematics* 123 (1986), 269–300.
- [Fo] G.B. Folland, Real analysis, modern techniques and their applications, *John Wiley and Sons* (1984).
- [HS] U. Haagerup, H. Schultz, Invariant Subspaces of Operators in a General II_1 -factor. Work in progress.
- [KR] R. V. Kadison, J. R. Ringrose, Fundamentals of the theory of operator algebras, Volume I, *Academic Press*, 1983.
- [Ku] K. Kuratowski, Topology, Volume I, 2nd. ed., Academic Press, London (1966).
- [Ne] E. Nelson, Notes on Non-commutative Integration, *Journal of Functional Analysis* 15 (1974), 103–116.